



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## Evaluating the implementation of the Mayo-Portland adaptability inventory-4 (MPAI-4) in three rehabilitation settings in Quebec: A mixed-methods study protocol

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2022-068866
Article Type:	Protocol
Date Submitted by the Author:	04-Oct-2022
Complete List of Authors:	<p>KENGNE TALLA, Pascaline; McGill University Faculty of Dentistry; Centre de recherche interdisciplinaire en readaptation du Montreal metropolitain</p> <p>Thomas, Alik; McGill University, School of Physical and Occupational Therapy, McGill University, Montréal, Canada; McGill University Faculty of Education</p> <p>Ataman, Rebecca; McGill University, School of Physical and Occupational Therapy, McGill University, Montréal, Canada; Centre de recherche interdisciplinaire en readaptation du Montreal metropolitain</p> <p>Auger, Claudine; Universite de Montreal Faculte de medecine, École de la réadaptation; Centre de recherche interdisciplinaire en readaptation du Montreal metropolitain</p> <p>McKerral, Michelle; Université de Montréal; Centre de recherche interdisciplinaire en readaptation du Montreal metropolitain</p> <p>Wittich, Walter; Universite de Montreal, School of Optometry</p> <p>Poncet, Frédérique; Université de Montréal, School of optometry; Centre de recherche interdisciplinaire en readaptation du Montreal metropolitain, Research</p> <p>Ahmed, Sara; McGill University, School of Physical and Occupational Therapy; Centre de recherche interdisciplinaire en readaptation du Montreal metropolitain</p>
Keywords:	STROKE MEDICINE, STATISTICS & RESEARCH METHODS, REHABILITATION MEDICINE, QUALITATIVE RESEARCH

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Evaluating the implementation of the Mayo-Portland adaptability inventory-4 (MPAI-4) in three rehabilitation settings in Quebec: A mixed-methods study protocol**

**Authors:** Kengne Talla, Pascaline<sup>1,2,3,4</sup>, Thomas, Aliko<sup>1,4,5,6</sup>, Ataman, Rebecca<sup>1,2,4,5</sup>, Auger, Claudine<sup>3,4,7</sup>, McKerral, Michelle<sup>3,4,8</sup>, Wittich, Walter<sup>2,4,9,10</sup>, Poncet, Frédérique<sup>1,2,4,9,10</sup>, Ahmed, Sara\*. <sup>1,2,3</sup>

Correspondence: Sara Ahmed: [sara.ahmed@mcgill.ca](mailto:sara.ahmed@mcgill.ca)

School of Physical and Occupational Therapy, McGill University, Montréal, Canada  
Integrated University Health and Social Services Centre for West-Central Montreal (CIUSSS du Centre-Ouest-de-l'Île-de-Montréal)  
Integrated University Health and Social Services Centre for South-Central Montreal (Institut universitaire sur la réadaptation en déficience physique de Montréal, CIUSSS du Centre-Sud-de-l'Île-de-Montréal)

**Keywords:** Implementation, rehabilitation, Mayo-Portland adaptability inventory (MPAI-4), integrated knowledge translation, mixed studies

**Abstract:**

**Introduction:** Stroke is a leading cause of morbidity and mortality worldwide, placing an immense burden on patients and the health system. Timely access to rehabilitation services and the use of standardized outcome measures is endorsed for optimizing patient rehabilitation outcomes and improving clinical decision-making. This project results from a provincially mandated recommendation to use the fourth version of the Mayo-Portland Adaptability Inventory (MPAI-4) to measure changes in social participation of stroke survivors and to maintain commitment to evidence-informed practices in stroke care. This protocol outlines the implementation process of the MPAI-4 for three rehabilitation

centers. The objectives are to: i) describe the context of MPAI-4 implementation; ii) determine clinical teams' readiness for change; iii) identify barriers and enablers to implementing the MPAI-4 and match the implementation strategies; iv) evaluate the MPAI implementation outcomes including the degree of integration of the MPAI-4 into clinical practice, and v) explore participants' experiences using the MPAI-4.

**Methods and analysis:** We will use a multiple case study design within an integrated knowledge translation approach with active engagement from key informants. Each case is a rehabilitation centre implementing MPAI-4. We will collect data from clinicians and program managers using mixed methods guided by several theoretical frameworks. Data sources include surveys, focus groups and patient charts. We will conduct descriptive, correlational, and content analyses. Ultimately, we will analyze, integrate data from qualitative and quantitative components and report them within and across participating sites. Results will provide insights about integrated knowledge translation within stroke rehabilitation settings that could be applied to future research projects.

**Ethics and dissemination:** The project received Institutional Review Board approval from the Centre for Interdisciplinary Research in Rehabilitation of Greater Montreal. We will disseminate results in peer-reviewed publications and at local, national, and international scientific conferences.

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- We will conduct this study in three rehabilitation centers in Canada using an integrated knowledge translation approach with clinicians and managers to better understand the implementation context thereby optimising likelihood of implementation success.
- The identification of barriers and enablers of successful implementation of evidence-informed practices in rehabilitation settings may help strengthen rehabilitation service planning.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

For peer review only

For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>

## BACKGROUND

Stroke is one of the main causes of morbidity and mortality [1-4], affecting an estimated 11.9 million people and accounting for 4.4 million deaths worldwide in 2017 [5-7]. Stroke survivors experience sequelae [8], including depression [9], loss of motor function [10], and vision loss [11]. Health and social consequences negatively influence the quality of life of stroke survivors [12], placing a burden on family and friends when they return home [13], and resulting in an economic burden to society [12].

Given the incidence of stroke and its impacts, a growing body of evidence suggests timely access to rehabilitation services improves stroke symptoms, patients' well-being [14-16], functional independence and social participation [17-19]. There is also mounting evidence to support the clinical use of standardized outcome measures [20, 21] to support the improvement of individuals' function and participation [22, 23], enhance patient rehabilitation outcomes [24-26], facilitate a patient-centered approach [27], and maintain clinical excellence and commitment to evidence-informed practices [28, 29]. An outcome measure used for these purposes is the Mayo-Portland Adaptability Inventory - version 4 (MPAI-4) which can assess both inpatient and outpatient rehabilitation patients' functional abilities and status [30-33]. Worldwide government health authorities and organizations have incorporated the MPAI-4 as part of their recommended practices in rehabilitation care for survivors of acquired brain injury. For instance, the National Outcome Info Database (USA) [13] and The Quebec Ministry of Health (*Ministère de la Santé et des Services sociaux*; MSSS) [34] have mandated the use of MPAI-4 in their local contexts. However, as with many new practices, the implementation of the MPAI-4 in clinical settings can be complex, multi-level, and thus, difficult to achieve. The implementation strategies that are targeted to the local context [35-37] may help to promote the adoption of evidence-informed practices [38], to improve patient and provider experiences related to the quadruple aim framework [39, 40] and ultimately, to inform the implementation success.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

This paper describes the protocol for a study that aims to evaluate the process of implementing the MPAI-4 using an integrated knowledge translation (iKT) approach [41], and evaluate its success (outcomes and impacts) in three stroke outpatient rehabilitation settings. IKT involves the engagement of stakeholders including managers and clinicians in each participating site in all the steps of the research process including the development of the research questions, selection of the study design and methodology, selection of outcome measures, data collection process, interpretation of the findings, dissemination of results [41, 42].

**The specific objectives are to:**

1. Describe the context in which each stroke rehabilitation site will implement the MPAI-4, and the potential strategies to improve implementation success.
2. Determine clinicians’ readiness to adopt the MPAI-4 in each site and across stroke rehabilitation sites.
3. Identify barriers and enablers to implementing the MPAI-4 within and across the stroke rehabilitation sites, as well as select and tailor the implementation strategies.
4. Evaluate the MPAI-4 implementation outcomes (acceptability, appropriateness, feasibility, adoption, and fidelity), including the degree to which the MPAI-4 is integrated into routine clinical practice within and across sites.
5. Explore clinicians’ and managers’ experiences of using MPAI-4 within and across sites.

## METHODS

### Study design

We will use a longitudinal descriptive multiple case study design [43] to comprehensively explore a phenomenon (i.e., the implementation of the MPAI-4) in its natural context [44]. According to Yin [45, 46], a case can be a decision, a program, an implementation process, an organizational change, a person, an event, or an entity that is context-dependent. In this study, a case will be a healthcare institution with its own stroke rehabilitation program. We will work with clinicians and managers to codevelop and execute the implementation plan. The use mixed-methods within the multiple case study [47, 48] will provide a deeper understanding of the factors influencing the clinicians' uptake of MPAI-4 while capturing the breadth of process and impact on outcomes.

### Implementation setting and description of the case

The study is a multi-centre project within the outpatient stroke rehabilitation programs in three regional health authorities in Quebec, Canada. Although these healthcare institutions (i.e., cases) share the same goals in terms of rehabilitation services offered to a stroke clientele, they differ in the territory and population density served, organizational culture and climate. Each of the rehabilitation programs includes between 20 and 45 clinicians, from different disciplines, providing services to between 200-300 outpatients annually, of various age groups, living with motor, language and/or sensory limitations.

### Description of the fourth version of the Mayo-Portland Adaptability Inventory (MPAI-4)

The MPAI-4 is freely available in many languages including French (Canadian) [30, 49-51] with good psychometric properties as the responsiveness [52-54], the cross-cultural validity and the reliability [49, 50]. The MPAI-4 includes 29 items classified in three main subscales: the ability, adjustment and participation [55]. All MPAI items are scored on a 5-point Likert scale (0 – 4) where 0 represents no



1  
2 limitations and 4 represents a severe issue interfering with activities more than 75% of the time. For  
3  
4 interpretation, this Likert scale must be converted to T-scores, with higher T-scores indicating lower  
5  
6 levels of functioning [30, 49-51].  
7  
8  
9

10  
11 **Patient and Public Involvement:** No patient involved in the project due to the COVID-10 pandemic  
12  
13 and its challenges.  
14

15  
16 **Implementation process**  
17  
18 We will work with clinicians and managers to iteratively codevelop, adapt and execute the  
19  
20 implementation plan for three major phases: pre-implementation, implementation and sustainability [56].  
21  
22 In this protocol we only focus on the first two phases. Implementation strategies will be suggested by the  
23  
24 clinical teams from their perceived barriers and enablers. Additional strategies will be tailored based on  
25  
26 barriers and enablers identified in each site as informed by the Consolidated Framework for  
27  
28 Implementation Research (CFIR) [37, 57, 58], (Table 1), as well as by the Expert Recommendations for  
29  
30 Implementing Change (ERIC) taxonomy [38]. The ERIC taxonomy is a compilation of implementation  
31  
32 strategies aiming to support selection and reporting of strategies used to address the potential  
33  
34 determinants of implementation [38, 59]. A summary of design and implementation process is available  
35  
36 in figure 1.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1: Implementation process and strategies**


Steps	Implementation Process	Approaches	Implementation strategies
Pre-implementation	Description of the context/Needs assessment (IT resources, human and materials resources)	<p>Determine with the support of the managers all the members of local implementation committee</p> <p>Meeting with IT team</p> <p>Regular meetings with each local implementation committee</p> <p>Choice and Adaptation to the electronic database ACCESS for the stroke program' needs</p> <p>Number of clinicians and patients in each clinical program</p> <p>Adaptations of the administration protocol of MPAI-4</p> <p>Choice of participating programs in each site</p>	<ul style="list-style-type: none"> <li>Facilitation strategies with an external facilitator who is a postdoctoral fellow with a great experience in knowledge translation and working in the clinical environments with various stakeholders including researchers, managers and patients. Another person with a facilitation role is an internal facilitator, member of the clinical team and of the local implementation committee, who works closely with the manager</li> <li>Local implementation committee composed of managers and clinical coordinators</li> <li>Inter-site implementation committee composed of managers and administrators</li> <li>Coach from traumatic brain injury (TBI) program</li> <li>Readiness survey</li> <li>Regular meetings with local implementation committee at each site (n=2 at least for all sites) to gain insights into the potential barriers and facilitators to implementing the MPAI-4, the technological aspects, the usability of MPAI-4</li> <li>Early implementation strategies as training, facilitation</li> <li>Set up booster training sessions every three months for people who need and for the new members of the clinical team</li> </ul>
	Training	Develop and adaptation of the training materials to each site (benefits, advantages, clinical utility, use, scoring and interpretation of MPAI-4 scores)	

Set up the training for instance  
virtual or in-person

**Table 1: Implementation process and strategies - Continued**

Steps	Implementation process	Approaches	Implementation strategies
Implementation phase	Pilot	Adaptation to each site for the duration of the pilot step and expectations during this step for instance 2 completed MPAI by clinicians where we will have one electronic and another one in - person	<ul style="list-style-type: none"><li>• Adaptation to the administration protocol</li><li>• Strategies to address barriers and enablers</li><li>• Regular meetings with implementation committees</li><li>• Sustainability strategies with the committees</li></ul>
	Full Scale	Choice of the date and duration of this step Define the expectations during this step for instance all clinicians will use the electronic version of MPAI Adaptation of the administration protocol after the pilot	<ul style="list-style-type: none"><li>• Adaptation to the administration protocol after the pilot and various questions from clinical team</li></ul>

**Figure 1: Implementation design and Timeline of MPAI-4 implementation project**



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Theoretical frameworks**

Three theoretical frameworks will be used to guide the pre-implementation and implementation phases. The first framework is the CFIR [57, 58] which we will use it to identify barriers and facilitators. The CFIR is composed of five domains of context: characteristics of the innovation, inner setting, outer setting, the processes that influence the implementation of innovations and integration into clinical routines, and characteristics of individuals [57]. We will adapt questions from the interview guide tool proposed by Damschroder and colleagues (2009) [60] .

The second framework is Proctor et al.’s Implementation Outcomes Framework (IOF) [61]. It proposes a taxonomy of implementation outcomes, patient outcomes and health service outcomes. In the case of the present project, we will focus on selected implementation outcomes, including acceptability, appropriateness, adoption, feasibility, and fidelity.

Lastly, we will use the Normalization Process Theory (NPT) to explore how the MPAI-4 becomes embedded and integrated into usual practice [62-64]. NPT theorises on the ways that people make sense of the work for implementing a complex intervention [62, 64]. In alignment with NPT, four constructs play a central role in generating the work of implementation: coherence/sense-making, cognitive participation, collective action and reflexive monitoring items [62, 64]. We will use the NPT to: i) assess the progress toward normalization over time and ii) compare the normalization (progress or outcomes) between sites in an implementation project [65].

**Pre-implementation phase**

This phase will consist of a local needs assessment to elicit information on available resources in each site (i.e., each case), clinicians’ readiness to adopt the MPAI-4 and the relevant implementation strategies

that may aid in activating the clinicians' adoption of the MPAI-4 and, ultimately, the implementation outcomes (aims 1 & 2) [66]. This phase is informed by the two first theoretical frameworks such as CFIR and Implementation outcomes framework.

**Design:** A mixed method (Qual-Quant) [47, 48] design will be used in this phase. The qualitative descriptive approach is used to understand the stakeholders' perspectives of their local context [46], and to conduct an in-depth needs assessment of available resources in each site. The organizational and clinical teams' readiness will be evaluated with the quantitative component.

**Participants:** We will form two sets of committees: one local implementation committee composed of managers and clinicians/clinical coordinators at each of the three sites; and one inter-site implementation committee composed of managers/administrators from all participating sites. For the qualitative component, we will invite all members of the local implementation committees and members of the information technology (IT) service. For the quantitative component, we will recruit all clinicians from all professions in rehabilitation for instance occupational therapists, physical therapists, speech language pathologists...etc.

**Data collection:** Measure of readiness will be obtained from two sources: i) focus groups with clinical and information technology teams; and ii) survey including two questionnaires for clinicians. The managers of the respective rehabilitation programs will contact the participants via email.

**Focus groups:** In the focus group [67, 68], we will collect information about the local context including the composition of the clinical team and clinical process, clinicians' and managers' knowledge of the MPAI-4, their needs, expectations and goals, the available resources, the process of data collection, and their unique organizational challenges (objective 1). We will collect information on the potential implementation strategies to increase the clinicians' buy-in and adoption of MPAI-4 from their own perspectives. Focus groups with information technology teams will be used to understand the existing

1 patient care software systems, IT resources, and the IT requirement to improve the implementation  
2 success.  
3  
4  
5

6 Two members of the research team (PKT and RA) will conduct a 60-minute focus group in person or  
7 virtually at each participating sites with members of the local implementation committee and IT  
8 members. We will develop the interview guide based on existing literature on MPAl-4 implementation  
9 [23, 30, 69, 70], the team’s experience in this content area, and questions from the CFIR interview guide  
10 tool [57, 60]. Interviews will be audio-recorded and transcribed verbatim.  
11  
12  
13  
14  
15  
16  
17

18  
19  
20 **Survey:** Quantitative data will be collected from a survey sent to all clinicians. The approximately 15  
21 minutes survey will be composed of questions related to two measurement tools such as the  
22 organizational Readiness for Implementing Change (ORIC) and the early implementation outcomes  
23 described below. Sociodemographic variables will be collected for instance gender, age group, site of  
24 work, profession. The survey will take approximately 10 minutes to fill out each measure tool for a  
25 total of 20 minutes.  
26  
27  
28  
29  
30  
31  
32  
33

34  
35  
36 ***Organizational Readiness for Implementing Change***  
37

38 The ORIC is based on Weiner’s organizational theory [71] and aims to evaluate the organization’s  
39 readiness to implement change including its commitment to change and its change efficacy as perceived  
40 by its members, and to guide them in the identification of strategies and resources relevant to their context  
41 [72]. The survey will contain 10-items scored on a 5-point Likert scale, ranging from “Disagree” to  
42 “Agree” [73]. It has been translated in many languages including Canadian French where the validation  
43 process has been conducted in a rehabilitation setting [73]. French and English versions of this tool have  
44 good psychometric properties including content validity, construct validity, and reliability [73].  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

***Implementation outcomes: Acceptability of Intervention Measure (AIM), Intervention Appropriateness Measure (IAM), and Feasibility of Intervention Measure (FIM))***

There are three briefs, validated, pragmatic and reliable measures developed by Weiner and collaborators and related to Proctor et al.'s IOF [61, 74]. Each measure is composed of four items per construct with ordinal five response options (from “completely disagree” to “completely agree”), giving a total of 12 questions [74]. Cut-off scores for interpretation not yet available; however, higher scores indicate greater acceptability, appropriateness, or feasibility [74]. There is support for AIM, IAM and FIM psychometric properties including good inter-item reliability and test-retest reliability [74]. We will use the original English versions and a non-validated French translation of these measures because there is no French version of this tool.

**Data Analysis:** Two members of the research team (PKT and RA) will conduct a content analysis of the qualitative data [75] using N\*Vivo software [76] and a quantitative data analysis using SAS version 12.1 [77]. They will anonymize the transcripts and review them for accuracy. Analysis will involve three phases [75]: 1) familiarize with the data; 2) organize the data with a categorization matrix; and, 3) report the data with the presentation of the described contents (meanings) of the categories and themes. Throughout this process, PKT and RA will meet regularly with one another and with the larger research team to discuss coding and potential recurring ideas and generate categories and themes as a part of a reflexive cycle. Any discrepancies will be resolved by discussion. Trustworthiness [78-80] will be supported through triangulation of data sources and collection methods (focus groups, meetings minutes, facilitators' notes, and surveys), using of multiple coders and several sites. Descriptive analyses of the quantitative data (e.g., mean, standard deviation, percentage) will be conducted to address variation in clinician readiness. We will explore the difference between sociodemographic variables of clinician readiness within and across sites. Qualitative and quantitative data will be integrated to further inform



1 the analysis as well as during the interpretation phase. We will use quotes to illustrate the most relevant  
2 findings from the qualitative data. We will generate tables with information on socio-demographic  
3 characteristics and from measurement tools.  
4  
5  
6  
7  
8  
9

10  
11 **Anticipated Results:** Results from this mixed-methods within the multiple case study will help to: 1)  
12 shed light on clinicians’ perceptions regarding the MPAI-4 and its compatibility with the clinical  
13 practice; and 2) inform and build a multicomponent implementation blueprint or plan of necessary  
14 resources tailored to the local contexts and to improve the implementation success. The identification of  
15 barriers and enablers informed by CFIR will guide the tailoring of the implementation strategies [59].  
16 We will consider implementation strategies from clinical teams as well from ERIC after matching with  
17 identified barriers and enablers. These strategies will be ranked and prioritize by the research team and  
18 the local implementation committees, for strategies most likely to increase the clinicians’ adoption of the  
19 MPAI-4 and to inform the next steps. The variation across the sites is expected to result in different  
20 adaptations to the MPAI-4 and/or implementation strategies to improve implementation success.  
21 Building on existing MPAI-4 materials used during previous implementation the MPAI-4 in traumatic  
22 brain injury rehabilitation settings [30, 51], we will develop the administration protocol of the MPAI-4  
23 in each participating site (who, what, when, where and how).  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 **Implementation phase**

44  
45 This phase will include two components: 1) a pilot period when each clinician administers the MPAI-4  
46 to four/six patients per site over a time selected by each site, followed by adaptations based on clinicians’  
47 and managers’ feedback; and 2) the full-scale implementation of the MPAI-4 across all sites (aims 3, 4  
48 & 5). We will identify the barriers and enablers to the uptake of the MPAI-4 by clinical teams and  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 evaluate the implementation outcomes. A summary of outcomes is available in Table 2. This phase is  
3  
4 informed by the three theoretical frameworks.  
5

6 **Design:** A qualitative descriptive approach will be used to identify barriers and enablers, and the  
7  
8 experiences of using MPAI-4. The level of integration of the MPAI-4 in the practices and the  
9  
10 implementation outcomes will be determined using a quantitative component.  
11  
12

13 **Participants:** All managers and clinicians from each site will be invited to participate via email. We aim  
14  
15 to recruit at least eight to ten clinicians, and one and/or two managers from each site to participate in  
16  
17 focus groups. We will invite all clinicians to fill out the survey.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 2: Process measures of implementation evaluation**

Process results and measures	Method of data collection	Data analysis
Description of barriers and enablers of MPAI-4' implementation	Meetings minutes from local needs assessment External and internal support field notes and site journal,  Focus groups with clinicians at 6 months (CFIR)	Content analysis
Description of the experiences with MPAI' implementation	Focus groups with clinicians and managers at 15 months	Content analysis
Description of the degree of normalization		Descriptive statistics
Feasibility:	Surveys (NOMAD) at 6, 12 and 15 months Access data	Quantitative: Descriptive statistics
1. Number of clinicians who used MPAI-4 and its subscales	Self-report by clinicians	Qualitative: content analysis
2. Number of clinicians who interpret MPAI-4 and its subscales	Field notes from external and internal support	
3. Time taken to complete MPAI-4 and all its subscales		
4. Time required to ACCESS/Redcap platform to a report		
5. Numbers of reports downloaded and put in the patient charts		
Fidelity:	Access data	Quantitative: Descriptive statistics
1. Number of missing subscales by clinicians	Field notes from internal and external support	Qualitative: content analysis
2. Number of missing case report forms		
3. Reasons for missing data		
Acceptability:	Surveys and self-report by clinicians	Quantitative: Descriptive statistics
1. % clinicians reporting MPAI-4 helpful in discussing symptoms/symptom management	Field notes form internal and external support	Qualitative: content analysis
2. % clinicians reporting ease of use and comprehensibility for MPAI-4 and technology systems		

**Table 2: Process measures of implementation evaluation - Co**

Outcomes measures	Method of data collection	Data analysis
<b>Appropriateness:</b> 1. % clinicians who find that MPAI-4 fit with workflow and clinical processes (e.g., MPAI-meaningful for clinical care, integrated in electronic health record system, linked clinical decision support) 2. MPAI-4 fit with clinic culture and values 3. The perceived relative advantage to use MPAI-4 (technology comfort, literacy level, meaningful for clinical condition)	Survey and self-reported by clinicians Field notes from external and internal support - PDSA	Quantitative: descriptive statistics Qualitative: content analysis
<b>Fidelity:</b> 1. Proportion of patients whose MPAI-4 was completed on expected times (admission, discharge, both) 2. % of targeted patients whose MPAI-4 was completed at the first individual intervention plan % of targeted patients whose MPAI was totally completed at discharge 3. Proportion of patients whose MPAI-4 scores were sent to the follow-up team, e.g., return-to-work program or CLSC 4. Proportion of targeted patients for whom the MPAI-4 was completed at discharge, so the total score decreased (improvement).	Access data	Quantitative: descriptive analysis
<b>Adoption:</b> Number of clinicians who used MPAI-4 over time (9, 12 and 15 months post training) Experiences/satisfaction/success MPAI- inform the PII same it is not discussed with the patient	Access data   Focus groups Field notes	Quantitative: descriptive analytics   Qualitative: content analysis

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Data collection:** Data collection will occur at three timepoints: at 6-, 12-, and 15- months following the pilot period defined in each site, during the full-scale implementation phase. The collected data will include information about: i) barriers and enablers to use of MPAI-4; ii) integration of MPAI-4 into clinical practice; iii) implementation outcomes (acceptability, appropriateness, feasibility, fidelity and adoption); and iv) experiences of the MPAI-4 use (for instance communication within and across sites, predefined goal as organizational, reflexivity, patient-centered care, patient-outcomes, technical and technological issues).

Focus group questions will be used to identify the factors that influence the uptake of the MPAI-4. We will conduct two focus groups of 90-minutes each [68]: a first to identify the barriers and enablers with the clinicians, and a second to explore managers' and clinicians' experiences of using MPAI-4. We will pilot-test the interview guide with one clinician and/or one manager to ensure the clarity of questions and their comprehensiveness. The first interview guide will be composed of open-ended questions addressing different constructs of the CFIR framework such as inner and outer settings, planning strategies, individual characteristics, intervention characteristics.

The second interview guide will consist of open-ended questions addressing clinicians and managers' viewpoints on their experiences with the use of MPAI-4 during this period, the strategies deployed for its implementation, the level of integration of the MPAI into the patient's treatment plan, their level of satisfaction and the lessons learned. Managers and clinicians will be recruited by email by a member of the research team. Before the focus group, participants will be asked to complete a brief sociodemographic questionnaire. We will use a similar approach for the focus group in the qualitative arm for the pre-implementation phase.

We will use the screening of MPAI-4 data from patient charts and three online, auto-administrated, and anonymous surveys to collect information on the MPAI-4 integration across time and the implementation measures using the Normalisation Measure Development questionnaire (*NoMAD*) [62, 81].

We will report MPAI-4 data from patient charts at different periods of administration (admission and discharge).

Underpinned by NPT [82, 83], the 20-item NoMAD assesses the normalisation of complex healthcare interventions such as the MPAI-4. It consists of three parts: Part A: sociodemographic information, Part B collecting information about the current use and likelihood of using the intervention in the future, and Part C comprising 20 items on 5-point Likert scale (1=completely agree to 5=completely disagree). This instrument has satisfactory psychometrics properties in French and English including the construct validity and the reliability [62]. Several implementation process outcomes will be collected from patients' charts. We will send the surveys to all clinicians in each site at different timelines. They will take approximatively 15 minutes to fill out it.

**Data analysis:** We will analyze the quantitative and qualitative data separately. The qualitative data will be used to inform the next steps in the implementation, mainly for choosing and tailoring implementation strategies. Both the quantitative and qualitative data will inform the last qualitative data collection on participants' experiences of the MPAI-4 use.

The qualitative data will be analyzed using a hybrid deductive-inductive approach [84]. Themes will be derived from the CFIR [57] while still allowing for emerging categories. Quantitative data from the NoMAD will be analyzed using descriptive analyses (e.g., mean, standard deviation, percentage). Internal consistency of all theoretical measures will be evaluated using Cronbach's alpha with an acceptable threshold defined as 0.70 [85]. We will estimate the MPAI-4 subscale scores as well as the

1  
2 overall score by data from patient charts. We will generate user progress reports by site, and across sites.  
3  
4 We will generate tables or figures with information on socio-demographic characteristics. We will use  
5  
6 quotes to illustrate the most relevant findings.  
7  
8  
9

10  
11 We will subsequently aggregate and integrate the data for a more comprehensive interpretation of the  
12  
13 results. Triangulation across various data sources theoretical applications, data collection and data  
14  
15 analysis techniques will be used to increase the depth of understanding of barriers and enablers to the  
16  
17 uptake the MPAI-4 in stroke rehabilitation settings. We will highlight all areas of convergence and  
18  
19 divergence across the participating rehabilitation sites.  
20  
21  
22

23  
24 **Anticipated Results:** This phase will generate a list of barriers and enablers, and potential tailored  
25  
26 implementation strategies matched to reduce the barriers and enhance and maintain the enablers [59].  
27  
28 The data will help to understand: the current use, the likelihood of using the MPAI-4 in the future,  
29  
30 clinicians' perceptions of the impact of MPAI-4 on their clinical practice, the perceived changes over the  
31  
32 time, and whether it could become a routine part of practice.  
33  
34  
35

36  
37  
38  
39 **DISCUSSION**  
40

41 We propose that successful implementation of the MPAI-4 requires an IKT approach with active  
42  
43 engagement from all stakeholders, including managers and clinicians in each of the participating clinical  
44  
45 rehabilitation programs. This study has the potential to contribute to the advancement of knowledge  
46  
47 regarding the implementation of the MPAI-4 by examining similarities and differences between sites,  
48  
49 highlighting the influence of context in the success of implementation (or not) that may influence  
50  
51 providers' experiences. The combination of multiple theoretical frameworks offers the opportunity to  
52  
53 map the broad barriers and enablers influencing clinicians' adoption of MPAI-4 uptake, and to select  
54  
55  
56  
57  
58  
59  
60

1 tailored strategies and to comprehensively measure the implementation processes and outcomes,  
2 including clinician's perceived clinical utility of MPAI-4 [69] as well as strategies that will increase the  
3 success of the MPAI-4's implementation in stroke rehabilitation programs.  
4  
5  
6  
7  
8  
9

10  
11 The implementation process of the MPAI-4 will provide us with data to facilitate communication  
12 between clinical teams across the care continuum. It will strengthen patient-centered practice and  
13 improve patient outcomes with the goal of optimizing social participation. This study will enhance the  
14 clinical utility of the MPAI-4 in stroke. Ultimately, we will conduct a comparison of our results on  
15 determinants with those of previous studies conducted with TBI patients [22, 30, 86]. Results of this  
16 study will provide insights about iKT approach within stroke rehabilitation settings that could be applied  
17 to future research projects.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

29 Despite the valuable insights that may be gained from this study, there are some limitations. Mixed  
30 methods with many data sources such as self-reported questionnaires, patient charts and focus groups  
31 may introduce response bias, that is, an under or overestimation of the expected behavior. To overcome  
32 this challenge, we will use triangulation across data sources, various theoretical applications, different  
33 methods of analysis and clinical teams' involvement to increase the in-depth understanding of our data  
34  
35  
36  
37  
38  
39  
40  
41  
42

### 43 **Ethics and dissemination**

44  
45 The project has received the Institutional Review Board (IRB) approval of CRIR (CRIR 1523-0221/MP-  
46 50-2022-968). Any protocol modifications will be submitted for approval to the IRB, prior to  
47 implementation. The project will be conducted from 2022 to 2024.  
48  
49  
50

51 The dissemination plan will be developed with the implementation team in each site. The iKT approach  
52 (participatory approach, reflective approach, local team) will allow for ongoing knowledge exchange  
53  
54  
55  
56  
57  
58  
59  
60



1 within and between participating sites. In addition, we will disseminate results in peer-reviewed  
2 publications and at local, national, and international scientific conferences/workshops. Reporting of this  
3 study will seek to satisfy the standards of Good Reporting of a Mixed Methods Study [87] and the  
4 Standards for Reporting Implementation Studies [88, 89].  
5  
6  
7  
8  
9

10  
11  
12  
13 **Acknowledgements:** We would like to acknowledge the leadership of managers in the participating  
14 rehabilitation centers, and we thank the members of various local implementation committees and all  
15 clinicians in these sites.  
16  
17  
18  
19

20  
21  
22 **Authors' contributions:** PKT led the writing of the manuscript. A research grant was obtained by SA  
23 and AT to support the proposed research. PKT, SA and AT produced the first draft of the article including  
24 all sections. CA, MM, WW, FP and RA contributed substantially to the study design and provided critical  
25 feedback on the manuscript. All authors read and approved the final version of the manuscript  
26  
27  
28  
29  
30

31  
32  
33 **Funding statement:** This study was supported by the *Pôle Universitaire en Réadaptation* (PUR) of the  
34 CRIR in 2018-2019 (N/A for grant number), and salary awards from the Fonds de recherche du Québec  
35 – Santé (FRQS) for SA, AT, CA, WW. This funding has been used to give students awards (PKT and  
36 RA). The implementation of the MPAI-4 is supported by the Biomedical Research and Informatics  
37 Living Laboratory for Innovative Advances of New Technologies (BRILLIANT) in Community  
38 Mobility Rehabilitation program funded by the Canadian Foundation of Innovation and the Ministry of  
39 Health of Quebec (#36053). CRIR is funded by the Fonds de recherche du Québec – Santé, et Société et  
40 culture.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

51  
52  
53  
54 **Competing interests' statement:** The authors declare no competing interests.  
55  
56  
57  
58  
59  
60

**Authors' information:**

1. School of Physical and Occupational Therapy, McGill University, Montréal, Canada
2. Integrated University Health and Social Services Centre for West-Central Montreal (CIUSSS du Centre-Ouest-de-l'Île-de-Montréal)
3. Integrated University Health and Social Services Centre for South-Central Montreal (Institut universitaire sur la réadaptation en déficience physique de Montréal, CIUSSS du Centre-Sud-de-l'Île-de-Montréal)
4. Centre for Interdisciplinary Research in Rehabilitation of Greater Montreal
5. Integrated Health and Social Services Centre of Laval (CISSS de Laval)
6. Institute of Health Sciences Education, McGill University, Montréal, Canada
7. School of Rehabilitation, Université de Montréal, Montréal, Canada
8. Department of Psychology, Université de Montréal, Montréal, Canada
9. School of Optometry, Université de Montréal, Montréal, Canada
10. Institut Nazareth et Louis-Braille, Integrated Health and Social Services Centre of Montérégie-Centre (CISSS de la Montérégie-Centre)

REFERENCES

1. Lindsay MP, Norving B, Sacco RL, Brainin M, Hacke W, Martins S, Pandian J, Feigin V. World Stroke Organization (WSO): Global Stroke Fact Sheet 2019. *Int J Stroke* 2019;14(8):806-17.

2. World stroke organization (WSO). Global Stroke Fact Sheet. 2019 [cited 2020 december, 10th]; Available from: [https://www.world-stroke.org/assets/downloads/WSO\\_Global\\_Stroke\\_Fact\\_Sheet.pdf](https://www.world-stroke.org/assets/downloads/WSO_Global_Stroke_Fact_Sheet.pdf).

3. Jolliffe L, Lannin NA, Cadilhac DA, Hoffmann T. Systematic review of clinical practice guidelines to identify recommendations for rehabilitation after stroke and other acquired brain injuries. *BMJ Open* 2018;8(2):e018791.

4. Johnson CO, Nguyen M, Roth GA, Nichols E, Alam T, Abate D et al. Stroke Collaborators. Global, regional, and national burden of stroke, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet Neurol* 2019;18(5):439-58.

5. Avan A, Digaleh H, Di Napoli M, Stranges S, Behrouz R, Shojaeianbabaei G, et al. Socioeconomic status and stroke incidence, prevalence, mortality, and worldwide burden: an ecological analysis from the Global Burden of Disease Study. *BMC Med* 2019; 17: 191-221.

6. Stanaway DJ, Afshin A, Gakidou E, Lim SS, Abate D, Abate KH et al. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study. *The Lancet* 2017; 392 (10159): 1923 – 94.

7. Institute for Health Metrics and Evaluation (IHME), Findings from the Global Burden of Disease Study 2017. Seattle, WA: IHME. 2018. p. 1-27.

8. Health and Stroke Foundation Canadian partnerships for stroke recovery, Background concepts in stroke rehabilitation. 2018: Canada. p. 1-34.

9. Ferro JM, Caeiro L, and Figueira ML. Neuropsychiatric sequelae of stroke. *Nat Rev Neurol* 2016; 12(5):269-80.
10. Zorowitz RD, Gillard PJ, and Brainin M. Poststroke spasticity: sequelae and burden on stroke survivors and caregivers. *Neurology* 2013; 80(3 Suppl 2): S45-52.
11. Rowe FJ, Hepworth LR, Howard C, Hanna KL, Cheyne CP, Currie J. High incidence and prevalence of visual problems after acute stroke: An epidemiology study with implications for service delivery. *PLoS One* 2019;14(3):e0213035.
12. Fondation des maladies du coeur et de l'AVC, Recommandations canadiennes pour les pratiques optimales de soins de l'AVC. Réadaptation, rétablissement et participation communautaire après un AVC: Deuxième partie : Les transitions et la participation communautaire après un AVC 2019; 1-82.
13. Ministère de la Santé et des Services sociaux (MSSS), Continuum de services pour les personnes à risque de subir ou ayant subi un accident vasculaire cérébral: Plan de mise en œuvre Phase 2016-2018, 2016; 1-64.
14. Cieza A. Rehabilitation the Health Strategy of the 21st Century, Really? *Arch Phys Med Rehabil* 2019; 100(11): 2212-14.
15. Negrini S, Mills JA, Arienti C, Kiekens C, Cieza A. "Rehabilitation Research Framework for Patients With COVID-19" Defined by Cochrane Rehabilitation and the World Health Organization Rehabilitation Programme. *Arch Phys Med Rehabil* 2021;102(7):1424-30.
16. Krug, E. and A. Cieza, Strengthening health systems to provide rehabilitation services. *Bulletin of the World Health Organization* 2017;95(3): 167.

17. Bernhardt J, Hayward KS, Kwakkel G, Ward NS, Wolf SL, Borschmann K, et al. Agreed definitions and a shared vision for new standards in stroke recovery research: The Stroke Recovery and Rehabilitation Roundtable taskforce. *Int J Stroke* 2017;12(5):444-50.

18. Pollock A, Baer G, Campbell P, Choo PL, Forster A, Morris J, et al. Physical rehabilitation approaches for the recovery of function and mobility following stroke. *Cochrane Database Syst Rev* 2014;(4):CD001920.

19. World Health Organization (WHO). Rehabilitation 2030 Initiative <https://www.who.int/initiatives/rehabilitation-2030>. 2017 [cited 2021 July, 20th].

20. Fricke, J., Measuring Outcomes in Rehabilitation: A Review. *Br J of Occupat Ther* 1993;56: 217 - 21.

21. Harris K, Dawson J, Gibbons E, Lim CR, Beard DJ, Fitzpatrick R, et al. Systematic review of measurement properties of patient-reported outcome measures used in patients undergoing hip and knee arthroplasty. *Patient Relat Outcome Meas*. 2016 Jul 25;7:101-8.

22. Institut national d'excellence en santé et en services sociaux (INESSS), Trousse de soutien à l'implantation clinique du MPAI-4. 2019: Québec, Québec. p. 1-46.

23. Malec JF. The Mayo-Portland Adaptability Inventory. The Center for Outcomes Measurement in Brain Injury. <http://www.tbims.org/combi/mpai>. 2005 [cited 2021 January 3rd].

24. Colquhoun HL, Lamontagne ME, Duncan EA, Fiander M, Champagne C, Grimshaw JM. A systematic review of interventions to increase the use of standardized outcome measures by rehabilitation professionals. *Clin Rehabil* 2017;31(3):299-309.

25. Sullivan JE, Crowner BE, Kluding PM, Nichols D, Rose DK, Yoshida R, Pinto Zipp G. Outcome measures for individuals with stroke: process and recommendations from the

- American Physical Therapy Association neurology section task force. *Phys Ther* 2013;93(10):1383-96.
26. Winstein CJ, Stein J, Arena R, Bates B, Cherney LR, Cramer SC, et al. American Heart Association Stroke Council, Council on Cardiovascular and Stroke Nursing, Council on Clinical Cardiology, and Council on Quality of Care and Outcomes Research. Guidelines for Adult Stroke Rehabilitation and Recovery: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association 2016;47(6):e98-e169.
27. Bellon K, Malec JF, Kolakowsky-Hayner SA. Mayo-Portland Adaptability Inventory-4. *J Head Trauma Rehabil* 2012; 27(4):314-6.
28. Accreditation Canada. Achieving Distinction in Stroke Care. 2019 [cited 2020 January, 7th]; Available from: <https://accreditation.ca/news/distinction-in-stroke-care/>.
29. Lageman SK. Commission on Accreditation of Rehabilitation Facilities (CARF), in *Encyclopedia of Clinical Neuropsychology* in: Kreutzer JS, DeLuca J, Caplan B. Editors. 2011, Springer New York: New York, NY. p. 647-50.
30. Guerrette MC, McKerral M. Validation of the Mayo-Portland Adaptability Inventory-4 (MPAI-4) and reference norms in a French-Canadian population with traumatic brain injury receiving rehabilitation. *Disabil Rehabil* 2021; 1-7.
31. Malec JF, Kragness M, Evans RW, Finlay KL, Kent A, Lezak MD. Further psychometric evaluation and revision of the Mayo-Portland Adaptability Inventory in a national sample. *J Head Trauma Rehabil* 2003;18(6):479-92.
32. Malec JF. Comparability of Mayo-Portland Adaptability Inventory ratings by staff, significant others and people with acquired brain injury. *Brain Injury* 2004;18(6): 563-75.
33. Williams E, Jackson H, Wagland J, Martini A. Community Rehabilitation Outcomes

for Different Stroke Diagnoses: An Observational Cohort Study. Arch Rehabil Res Clin Transl 2020;2(2):100047.

34. Ministère de la Santé et des Services socaux (MSSS). Continuum de services pour les personnes à risque de subir ou ayant subi un accident vasculaire cérébral, Quebec, 2018: 1-89.

35. Harvey G, Loftus-Hills A, Rycroft-Malone J, Titchen A, Kitson A, McCormack B, et al. Getting evidence into practice: the role and function of facilitation. J Adv Nurs 2002;37(6):577-88.

36. Kitson AG, Harvey G, McCormack B. Enabling the implementation of evidence based practice: a conceptual framework. Quality in Health Care 1998;7(3): p. 149-158.

37. Damschroder LJ. Clarity out of chaos: Use of theory in implementation research. Psychiatry Research 2020; **283**:112461.

38. Powell BJ, Waltz TJ, Chinman MJ, Damshoreder LJ, Smith JL, Monica MM et al. A refined compilation of implementation strategies: results from the Expert Recommendations for Implementing Change (ERIC) project. Implement Sci 2015;10: 21.

39. Bodenheimer T, Sinsky C. From triple to quadruple aim: care of the patient requires care of the provider. Annals of family medicine 2014; 12(6): 573-76.

40. Sikka R, Morath JM, Leape L. The Quadruple Aim: care, health, cost and meaning in work. BMJ Qual Saf 2015;24(10):608-10.

41. Kothari A, McCutcheon C, Graham ID. Defining Integrated Knowledge Translation and Moving Forward: A Response to Recent Commentaries. Int J Health Policy Manag 2017; 6(5):299-300.

42. Graham ID, Kothari A., McCutcheon C, [On behalf of the Integrated Knowledge Translation Research Network Project Leads](#). Moving knowledge into action for more effective practice, programmes and policy: protocol for a research programme on integrated knowledge translation. *Implement Sci* 2018; 13: 22.
43. Thomas G. A Typology for the Case Study in Social Science following a review of definition, discourse, and structure. *Qual Inquiry* 2011; 17(6):511-21.
44. Eisenhardt KM, Graebner ME. Theory Building From Cases: Opportunities And Challenges. *Academy of Manag J* 2007; 50(1):25-32.
45. Yin RK. Case study research: Design and methods (4th Ed.). Thousand Oaks, CA: Sage. 2009.
46. Yin RK. Case Study Research and Applications: Design and Methods (6th ed.). Thousand Oaks, CA: Sage. 2018.
47. Creswell JW. Research Design: Qualitative, Quantitative, and mixed methods approaches. 2009, California: Sage Publication.
48. Palinkas LA, Aarons GA, Horwitz S, Chamberlain P, Hurlburt M, Landsverk J. Mixed method designs in implementation research. *Adm Policy Ment Health* 2011;38(1):44-53.
49. Hamed R, Tariah HA, Malkawi S, Holm MB. The Arabic version of the Mayo-Portland Adaptability Inventory 4: a validation study. *Int J Rehabil Res* 2012;35(3):243-7.
50. Cattalani R, Corsini D, Posteraro L, Agosti M, Saccavini M. The Italian version of the Mayo-Portland Adaptability Inventory-4. A new measure of brain injury outcome. *Eur J Phys Rehabil Med* 2009;45(4):513-9.



51. McKerral M, Léveillé G, Goulet P, Vincent P. Manuel MPAI-4 Version canadienne-française. The Center for Outcome Measurement in Brain Injury. Available from: <http://www.tbims.org/combi/mpai/>. 2014 [cited 2021 June 6th].

52. Jackson D, Seaman K, Sharp K, Singer R, Wagland J, Turner-Stokes L. Staged residential post-acute rehabilitation for adults following acquired brain injury: A comparison of functional gains rated on the UK Functional Assessment Measure (UK FIM+FAM) and the Mayo-Portland Adaptability Inventory (MPAI-4). *Brain Inj* 2017;31(11):1405-13.

53. Nayar M, Vanderstay R, Siegert RJ, Turner-Stokes L. The UK Functional Assessment Measure (UK FIM+FAM): Psychometric Evaluation in Patients Undergoing Specialist Rehabilitation following a Stroke from the National UK Clinical Dataset. *PLoS One* 2016;11(1):e0147288.

54. Altman IM, Swick S, Malec JF. Effectiveness of Home- and Community-Based Rehabilitation in a Large Cohort of Patients Disabled by Cerebrovascular Accident: Evidence of a Dose-Response Relationship. *Archives of Physical Medicine and Rehabilitation* 2013; 94(9): 1837-41.

55. Malec, J.F. The Mayo-Portland Adaptability Inventory. The Center for Outcome Measurement in Brain Injury. <http://www.tbims.org/combi/mpai>. 2005 [cited 2021 January 3rd].

56. Kilbourne AM, Goodrich DE, Miake-Lye I, Braganza MZ, Bowersox NW. Quality Enhancement Research Initiative Implementation Roadmap: Toward Sustainability of Evidence-based Practices in a Learning Health System. *Med Care* 2019;57 Suppl 10 Suppl 3(10 Suppl 3):S286-93.

57. Damschroder LJ, Aron DC, Keith RE, Kirsh SR, Alexander JA, Lowery JC. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci* 2009; 4(1): p. 50.
58. Esmail R, Hanson H, Holroyd-Leduc J, Brown, S, Striffler L, Straus SE, Niven DJ. A scoping review of full-spectrum knowledge translation theories, models, and frameworks. *Implement Sci* 2020;15:11.
59. Waltz TJ, Powell BJ, Fernández ME, Abadie B, Damschroder LJ. Choosing implementation strategies to address contextual barriers: diversity in recommendations and future directions. *Implement Sci* 2019;14(1):42.
60. Consolidated Framework for Implementation research (CFIR). Welcome to the Interview Guide Tool: CFIR Domains [https://cfirguide.org/guide/app/#/guide\\_select](https://cfirguide.org/guide/app/#/guide_select). [cited 2021 January, 12].
61. Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A et al. Outcomes for implementation research: conceptual distinctions, measurement challenges, and research agenda. *Adm Policy Ment Health* 2011;38(2):65-76.
62. Finch TL, Girling M, May CR, Mair FS, Murray E, Treweek S et al. Improving the normalization of complex interventions: part 2 - validation of the NoMAD instrument for assessing implementation work based on normalization process theory (NPT). *BMC Med Res Methodol* 2018;18(1):135.
63. Finch TL, Rapley T, Girling M, Mair FS, Murray E, Treweek S et al. Improving the normalization of complex interventions: measure development based on normalization process theory (NoMAD): study protocol. *Implement Sci* 2013;8-43.
64. Rapley T, Girling M, Mair FS, Murray E, Treweek S, McColl E, Steen IN, May CR,

Finch TL. Improving the normalization of complex interventions: part 1 - development of the NoMAD instrument for assessing implementation work based on normalization process theory (NPT). BMC Med Res Methodol 2018;18(1):133.

65. Vis C, Ruwaard J, Finch T, Rapley T, de Beurs D, van Stel H et al. Toward an Objective Assessment of Implementation Processes for Innovations in Health Care: Psychometric Evaluation of the Normalization Measure Development (NoMAD) Questionnaire Among Mental Health Care Professionals. J Med Internet Res 2019;21(2):e12376.

66. Lewis CC, Boyd MR, Walsh-Bailey C, Lyon AR, Beidas R, Mittman B.. A systematic review of empirical studies examining mechanisms of implementation in health. Implement Sci 2020; 15:21.

67. Carpenter C. Chap 5: Using qualitative focus groups to evaluate health programmes and service delivery in Qualitative research in evidence-based rehabilitation: Carpenter C (ed). 2004; 51-64.

68. Krueger RA, Casey MA. Focus groups : a practical guide for applied research. 2009: Thousand Oaks, California : SAGE.

69. Kean J, Malec JF, Cooper DB, Bowles AO. Utility of the Mayo-Portland adaptability inventory-4 for self-reported outcomes in a military sample with traumatic brain injury. Arch Phys Med Rehabil 2013;94(12):2417-24.

70. Wilde EA, Whiteneck GG, Bogner J, Bushnik T, Cifu DX, Dikmen S et al. R6commendations for the use of common outcome measures in traumatic brain injury research. Arch Phys Med Rehabil 2010;91(11):1650-1660.e17.

71. Weiner BJ. A theory of organizational readiness for change. Implement Sci 2009; 4(1):67.

72. Shea CM, Jacobs SR, Esserman DA, Bruce K, Weiner BJ. Organizational readiness for implementing change: a psychometric assessment of a new measure. *Implementation Sci* 2014;9:7.
73. Ruest M, Léonard G, Thomas A, Desrosiers J, Guay M. French cross-cultural adaptation of the Organizational Readiness for Implementing Change (ORIC). *BMC Health Serv Res* 2019;19:535.
74. Weiner BJ, Lewis CC, Stanick C, Powell BJ, Dorsey CN, Clary AS et al. Psychometric assessment of three newly developed implementation outcome measures. *Implement Sci* 2017;29;12(1):108.
75. Elo S, Kyngäs H. The qualitative content analysis process. *J Adv Nurs* 2008;62(1):107-15.
76. N\*Vivo Software in Qualitative Data Analysis: QSR International. 2020 [cited 2020 september 25th]; Available from: [https://www.qsrinternational.com/nvivo-qualitative-data-analysis-software/support-services/customer-hub/research-ready-\(1\)/research-ready](https://www.qsrinternational.com/nvivo-qualitative-data-analysis-software/support-services/customer-hub/research-ready-(1)/research-ready).
77. SAS Institute Inc, SAS. Version 12.1. <https://support.sas.com/documentation/onlinedoc/stat/121/intro.pdf>. 2020.
78. Shenton AK. Strategies for Ensuring Trustworthiness in Qualitative Research Projects. *Education for Information* 2004;22:63-75.
79. Graneheim UH, Lundman B. Qualitative content analysis in nursing research: concepts, procedures and measures to achieve trustworthiness. *Nurse Educ Today* 2004;24(2): 105-12.
80. Elo S, Kääriäinen M, Kanste O, Pölkki T, Utriainen K, Kyngäs, H. Qualitative

Content Analysis: A Focus on Trustworthiness. SAGE Open 2014;4:1-10.

81. ImpleMentAll. Normalization Measure Development Questionnaire (NoMAD), (2018, October 16). Retrieved from <http://www.implementall.eu/>. 2018 [cited 2021 June 25].

82. Finch, T.L., et al., *Nomad: Implementation measure based on Normalization Process Theory.[Measurement instrument]*. Retrieved from <http://www.normalizationprocess.org>. 2015.

83. May CR, Cummings A, Girling M, Bracker M, Mair FS, May CM et al. Using Normalization Process Theory in feasibility studies and process evaluations of complex healthcare interventions: a systematic review. Implement Sci 2018; 13:80.

84. Hsieh HF, Shannon S. Three Approaches to Qualitative Content Analysis. Qualitative health research 2005; 15: 1277-88.

85 . Cronbach LJ. Coefficient alpha and the internal structure of tests. Psychometrika,1951. 16: 297-334.

86. Lamontagne ME, Bussi res A, Kairy D, Ahmed S, Zidarov D, Messier F.  valuation des effets de l'implantation de quatre nouvelles pratiques cliniques fond es sur des donn es probantes- Phase 2 2017. p. 1-185. <https://ciusssmcq.ca/telechargement/633/evaluation-des-effets-de-l-implantation-de-quatre-nouvelles-pratiques-cliniques>

87. O'Cathain A, Murphy E, Nicholl J. The quality of mixed methods studies in health services research. J Health Serv Res Policy 2008;13(2):92-8.

88. Smith JD, LI DH, Rafferty MR. The Implementation Research Logic Model: a < method for planning, executing, reporting, and synthesizing implementation projects. Implement Sci 2020;15(1):84.

89. Hoffmann TC, Glaszion PP, Boutron I, Milne R, Perera R, Moher D et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. BMJ : Br Med J 2014;348: g1687.

For peer review only

# BMJ Open

## Evaluating the implementation of the Mayo-Portland adaptability inventory-4 (MPAI-4) in three rehabilitation settings in Quebec: A mixed-methods study protocol

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2022-068866.R1
Article Type:	Protocol
Date Submitted by the Author:	26-Apr-2023
Complete List of Authors:	KENGNE TALLA, Pascaline; McGill University Faculty of Dentistry; Centre de recherche interdisciplinaire en readaptation du Montreal metropolitain Thomas, Alik; McGill University, School of Physical and Occupational Therapy, McGill University, Montréal, Canada; McGill University Faculty of Education Ataman, Rebecca; McGill University, School of Physical and Occupational Therapy, McGill University, Montréal, Canada; Centre de recherche interdisciplinaire en readaptation du Montreal metropolitain Auger, Claudine; Universite de Montreal Faculte de medecine, École de la réadaptation; Centre de recherche interdisciplinaire en readaptation du Montreal metropolitain McKerral, Michelle; Université de Montréal; Centre de recherche interdisciplinaire en readaptation du Montreal metropolitain Wittich, Walter; Universite de Montreal, School of Optometry Poncet, Frédérique; Université de Montréal, School of optometry; Centre de recherche interdisciplinaire en readaptation du Montreal metropolitain, Research Ahmed, Sara; McGill University, School of Physical and Occupational Therapy; Centre de recherche interdisciplinaire en readaptation du Montreal metropolitain
<b>Primary Subject Heading</b>:	Rehabilitation medicine
Secondary Subject Heading:	Evidence based practice
Keywords:	STROKE MEDICINE, STATISTICS & RESEARCH METHODS, REHABILITATION MEDICINE, QUALITATIVE RESEARCH

SCHOLARONE™  
Manuscripts

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

**Evaluating the implementation of the Mayo-Portland adaptability inventory-4 (MPAI-4) in three rehabilitation settings in Quebec: A mixed-methods study protocol**

**Authors:** Kengne Talla, Pascaline<sup>1,2,3,4</sup>, Thomas, Aliko<sup>1,4,5,6</sup>, Ataman, Rebecca<sup>1,2,4,5</sup>, Auger. Claudine<sup>3,4,7</sup>, McKerral, Michelle<sup>3,4,8</sup>, Wittich, Walter<sup>2,4,9,10</sup>, Poncet, Frédérique<sup>1,2,4,9,10</sup>, Ahmed, Sara\*. <sup>1,2,3</sup>

Correspondence: Sara Ahmed: [sara.ahmed@mcgill.ca](mailto:sara.ahmed@mcgill.ca)

School of Physical and Occupational Therapy, McGill University, Montréal, Canada

Integrated University Health and Social Services Centre for West-Central Montreal (CIUSSS du Centre-Ouest-de-l'Île-de-Montréal)

Integrated University Health and Social Services Centre for South-Central Montreal (Institut universitaire sur la réadaptation en déficience physique de Montréal, CIUSSS du Centre-Sud-de-l'Île-de-Montréal)

Keywords: Implementation, rehabilitation, Mayo-Portland Adaptability Inventory (MPAI-4), integrated knowledge translation, mixed studies

1

For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>



## Abstract:

**Introduction:** Stroke is a leading cause of morbidity and mortality worldwide, placing an immense burden on patients and the health system. Timely access to rehabilitation services can improve stroke survivors' quality of life. The use of standardized outcome measures is endorsed for optimizing patient rehabilitation outcomes and improving clinical decision-making. This project results from a provincially mandated recommendation to use the fourth version of the Mayo-Portland Adaptability Inventory (MPAI-4) to measure changes in social participation of stroke survivors and to maintain commitment to evidence-informed practices in stroke care. This protocol outlines the implementation process of the MPAI-4 for three rehabilitation centers. The objectives are to: i) describe the context of MPAI-4 implementation; ii) determine clinical teams' readiness for change; iii) identify barriers and enablers to implementing the MPAI-4 and match the implementation strategies; iv) evaluate the MPAI-4 implementation outcomes including the degree of integration of the MPAI-4 into clinical practice, and v) explore participants' experiences using the MPAI-4.

**Methods and analysis:** We will use a multiple case study design within an integrated knowledge translation approach with active engagement from key informants. Each case is a rehabilitation centre implementing MPAI-4. We will collect data from clinicians and program managers using mixed methods guided by several theoretical frameworks. Data sources include surveys, focus groups and patient charts. We will conduct descriptive, correlational, and content analyses. Ultimately, we will analyze, integrate data from qualitative and quantitative components and report them within and across participating sites. Results will provide insights about integrated knowledge translation within stroke rehabilitation settings that could be applied to future research projects.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

**Ethics and dissemination:** The project received Institutional Review Board approval from the Centre for Interdisciplinary Research in Rehabilitation of Greater Montreal. We will disseminate results in peer-reviewed publications and at local, national, and international scientific conferences.

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- The study will use an integrated knowledge translation approach with clinicians and managers in rehabilitation centres.
  - The identification of the barriers and enablers for the successful implementation of MPAI-4 may help to better understand the implementation context.
  - The use of mixed-methods within the multiple case study will provide a deeper understanding of the factors influencing the clinicians' use of MPAI-4.
  - The triangulation of data collection and analysis methods guided by implementation science frameworks will help in optimizing implementation success.
  - The two data collection methods (i.e., self-reported surveys and focus groups) may introduce some bias.

## BACKGROUND

Stroke is one of the main causes of morbidity and mortality [1-4], affecting an estimated 11.9 million people, and accounting for 4.4 million deaths worldwide in 2017 [5-7]. Stroke survivors experience sequelae [8], including depression [9], loss of motor function [10], and vision loss [11]. Health and social consequences negatively influence the quality of life of stroke survivors [12], placing a burden on their family and friends when they return home [13], and resulting in an economic burden to society [12].

Given the incidence of stroke and its impacts, a growing body of evidence suggests timely access to rehabilitation services improves stroke symptoms, patients' well-being [14-16], functional independence and social participation [17-19]. There is also mounting evidence to support the clinical use of standardized outcome measures [20, 21] to support the improvement of individuals' function and participation [22, 23], enhance patient rehabilitation outcomes [24-26], facilitate a patient-centered approach [27], and maintain clinical excellence and commitment to evidence-informed practices [28, 29]. An outcome measure used for these purposes is the Mayo-Portland Adaptability Inventory - version 4 (MPAI-4) which can assess both inpatient and outpatient rehabilitation patients' functional abilities and status [30-33]. Worldwide government health authorities and organizations have incorporated the MPAI-4 as part of their recommended practices in rehabilitation care for survivors of acquired brain injury. For instance, the National Outcome Info Database (USA) [13] and the Quebec Ministry of Health (*Ministère de la Santé et des Services sociaux*; MSSS) [34] have mandated the use of MPAI-4 in their local contexts. However, as with many new practices, the implementation of the MPAI-4 in clinical settings can be complex, multi-level, and thus, difficult to achieve. The implementation strategies that are targeted to the local context [35-37] may help to promote the adoption of evidence-informed practices [38], to improve patient and provider experiences related to the Quadruple aim framework [39, 40] and ultimately, to inform the implementation success. In fact, the Quadruple aim framework describes the importance of health care improvements and transformation efforts of the health care system, including

1 improving the health of populations, patients' experience of care, health care providers' experience and  
2  
3  
4 reducing the cost of care with the intention of improving health equity.  
5  
6

7 This paper describes the protocol for a study that aims to evaluate the process of implementing the MPAI-  
8  
9 4 using an integrated knowledge translation (iKT) approach [41], and to evaluate its success (outcomes  
10  
11 and impacts) in three stroke outpatient rehabilitation settings. IKT involves the engagement of  
12  
13 stakeholders including managers and clinicians in each participating site in all the steps of the research  
14  
15 process including the development of the research questions, selection of the study design and  
16  
17 methodology, selection of the outcome measures, data collection process, interpretation of the findings,  
18  
19 and dissemination of the results [41, 42].  
20  
21  
22  
23  
24  
25  
26

27 **The specific objectives are to:**  
28  
29

- 30 1. Describe the context in which each stroke rehabilitation site will implement the MPAI-4, and the  
31  
32 potential strategies to improve implementation success.  
33  
34 2. Determine clinicians' readiness to adopt the MPAI-4 in each site and across stroke rehabilitation  
35  
36 sites.  
37  
38 3. Identify barriers and enablers to implementing the MPAI-4 within and across the stroke rehabilitation  
39  
40 sites, as well as select and tailor the implementation strategies.  
41  
42  
43 4. Evaluate the MPAI-4 implementation outcomes (acceptability, appropriateness, feasibility, adoption,  
44  
45 and fidelity), including the degree to which the MPAI-4 is integrated into routine clinical practice  
46  
47 within and across sites.  
48  
49

50 Explore clinicians' and managers' experiences of using MPAI-4 within and across sites.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## METHODS

### Study design

We will use a longitudinal descriptive multiple case study design [43] to comprehensively explore a phenomenon (i.e., the implementation of the MPAI-4) in its natural context [44]. According to Yin [45, 46], a case can be a decision, a program, an implementation process, an organizational change, a person, an event, or an entity that is context-dependent. In this study, a case will be a healthcare institution with its own stroke rehabilitation program. We will work with clinicians and managers to codevelop and execute the implementation plan. The use of mixed-methods within the multiple case study [47, 48] will provide a deeper understanding of the factors influencing clinicians' use of MPAI-4 while capturing the breadth of the process and the impact on outcomes.

### Implementation setting and description of the case

The study is a multi-centre project within the outpatient stroke rehabilitation programs in three regional health authorities in Quebec, Canada. Although these healthcare institutions (i.e., cases) share the same goals in terms of rehabilitation services offered to a stroke clientele, they differ in the territory and the population density served, and the organizational culture and climate. Each of the rehabilitation programs includes between 20 and 45 clinicians, from different disciplines, providing services to between 200-300 outpatients annually of various age groups, living with motor, language and/or sensory limitations.

### Description of the fourth version of the Mayo-Portland Adaptability Inventory (MPAI-4)

The MPAI-4 is freely available in many languages including French (Canadian) [30, 49-51], with generally high quality evidence of strong psychometric properties [52] including responsiveness (defined as the ability of the MPAI-4 to detect changes in a patient in rehabilitation over time) [53-55], cross-

1 cultural validity and reliability [49, 50]. The MPAI-4 includes 29 items classified in three main subscales:  
2 ability, adjustment and participation [56]. All MPAI-4 items are scored on a 5-point Likert scale (0 – 4)  
3 where 0 represents no limitations and 4 represents a severe issue interfering with activities more than  
4 75% of the time. For interpretation, this Likert scale must be converted to T-scores, with higher T-scores  
5 indicating lower levels of functioning [30, 49-51].  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16

17 **Patient/ Public Involvement:** There was no patient involvement in the project design and development  
18 of the protocol due to the COVID-19 pandemic and its challenges. However, the patients will be involved  
19 in the implementation project as they will be assessed using the MPAI-4 by the clinicians as part of  
20 routine rehabilitation care. There is no expected research data collection with patients on the  
21 implementation process and the use of MPAI-4 results in the clinical decision-making. However, the  
22 original funding included a budget for compensating patient partners.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

35 **Implementation process**

36 We will work with clinicians and managers to iteratively codevelop, adapt, and execute the  
37 implementation plan for three major phases: pre-implementation, implementation and sustainability [57].  
38 In this protocol we only focus on the first two phases. The study will last from June 2020 to December  
39 2023 (Figure 1). The implementation strategies will be suggested by the clinical teams based on their  
40 perceived barriers and enablers during the pre-implementation phase. Additional strategies will be  
41 tailored based on the barriers and enablers identified in each site as informed by the Consolidated  
42 Framework for Implementation Research (CFIR) [37, 58, 59], (Table 1), as well as by the Expert  
43 Recommendations for Implementing Change (ERIC) taxonomy [38] during the implementation phase.  
44 The ERIC taxonomy is a compilation of implementation strategies aiming to support selection and  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

reporting of strategies used to address the potential determinants of implementation [38, 60]. A summary of design and implementation process is available in figure 1.

For peer review only

**Table 1: Implementation process and strategies**

Steps	Implementation process	Approaches	Implementation strategies
Pre-implementation	Description of the context/Needs assessment (IT resources, human and materials resources)	<p>Involvement of managers and all the members of local implementation committee.</p> <p>Meeting with IT team</p> <p>Regular meetings with each local implementation committee</p> <p>Identification and adaptation to the electronic database</p> <p>Access for the stroke program’ needs</p> <p>Number of clinicians and patients in each clinical program</p> <p>Adaptations of the administration protocol of MPAI-4</p> <p>Identification of participation in programs in each site</p>	<ul style="list-style-type: none"><li>• Facilitation strategies with an external facilitator who is a postdoctoral fellow with extensive experience in knowledge translation and working in the clinical environment with various stakeholders including researchers, managers and patients (more than eight years working with the clinical teams). Another person with a facilitation role is an internal facilitator, member of the clinical team and of the local implementation committee, who works closely with the manager</li><li>• Local implementation committee composed of managers and clinical coordinators</li><li>• Inter-site implementation committee composed of managers and administrators</li><li>• Coach from traumatic brain injury (TBI) program</li><li>• Readiness survey</li><li>• Regular meetings with local implementation committee at each site (n=2 at least for all sites) to gain insights into the potential barriers and facilitators to implementing the MPAI-4, the technological aspects, the usability of MPAI-4</li><li>• Early implementation strategies as training, facilitation</li></ul>
	Training	<p>Development and adaptation of the training materials to each site (benefits, advantages, clinical utility, use, scoring and interpretation of MPAI-4 scores)</p>	<ul style="list-style-type: none"><li>• Set up booster training sessions every three months for people who need and for the new members of the clinical team</li></ul>



Set up the training for instance  
virtual or in-person

**Table 1: Implementation process and strategies - Continued**

Steps	Implementation process	Approaches	Implementation strategies
Implementation phase	Pilot	Adaptation to each site for the duration of the pilot step.  In-person or electronic administration of the MPAI-4	Adaptation to the administration protocol  Strategies to address barriers and enablers  Regular meetings with implementation committees  Sustainability strategies with the committees
	Main study	Defining the date and duration of this step  Defining the expectations during this step for instance all clinicians will use the electronic version of MPAI-4  Adaptation of the administration protocol after the pilot	Adaptation to the administration protocol after the pilot and various questions from clinical team

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Theoretical frameworks**

Three theoretical frameworks will be used to guide the pre-implementation and implementation phases. The first framework is the CFIR [58, 59] which we will use it to identify barriers and facilitators. The CFIR is composed of five domains of context: characteristics of the innovation, inner setting, outer setting, the processes that influence the implementation of innovations and integration into clinical routines, and characteristics of individuals [58]. We will adapt questions from the interview guide tool proposed by Damschroder and colleagues (2009) [61].

The second framework is Proctor et al.'s Implementation Outcomes Framework (IOF) [62]. It proposes a taxonomy of implementation outcomes, patient outcomes and health service outcomes. In the case of the present project, we will focus on selected implementation outcomes, including acceptability, appropriateness, adoption, feasibility, and fidelity.

Lastly, we will use the Normalization Process Theory (NPT) to explore how the MPAI-4 becomes embedded and integrated into usual practice [63-65]. NPT theorises on the ways that people make sense of the work for implementing a complex intervention [63, 65]. In alignment with NPT, four constructs play a central role in the process of implementation: coherence/sense-making, cognitive participation, collective action and reflexive monitoring [63, 65]. We will use NPT to: i) assess progress toward normalization over time and ii) compare normalization (progress or outcomes) between sites in an implementation project [66].

**Pre-implementation phase**

This phase will consist of a local needs assessment to elicit the information on the available resources in each site (i.e., each case), the clinicians' readiness to adopt the MPAI-4 and the relevant implementation strategies that may aid in activating the clinicians' adoption of the MPAI-4 and, ultimately, the implementation outcomes (objectives 1 & 2) [67]. This phase is informed by the two first theoretical frameworks, CFIR and the IOF.

**Design:** A mixed methods (Qual-Quant) design [47, 48] will be used in this phase. A qualitative descriptive approach will be used to understand the stakeholders' perspectives of their local context [46], and to conduct an in-depth needs assessment of available resources in each site. The organizational and clinical teams' readiness will be evaluated with the quantitative component.

**Participants:** We will form two committees: one local implementation committee composed of managers and clinicians/clinical coordinators at each of the three sites; and one inter-site implementation committee composed of managers/administrators from all participating sites. For the qualitative component, we will invite all members of the local implementation committees and members of the information technology (IT) services to participate. For the quantitative component, we will recruit clinicians from all professions in participating sites who can administer the MPAI-4 after its implementation, including for instance occupational therapists, physical therapists, psychologists, kinesiologists, social workers, and speech-language pathologists.

**Data collection:** Information on readiness will be obtained from two sources: i) focus groups with clinical and information technology teams; and ii) a survey including two questionnaires for clinicians. The managers of the respective rehabilitation programs will contact the participants via email.

**Focus groups:** In the focus group [68, 69], we will collect information about the local context including the composition of the clinical team and clinical process, clinicians’ and managers’ knowledge of the MPAI-4, their needs, expectations and goals, the available resources, the process of data collection, and their unique organizational challenges (objective 1). We will collect information on the potential implementation strategies to increase the clinicians’ buy-in and adoption of MPAI-4 from their own perspectives. Focus groups with information technology will be used to understand the existing patient care software systems, IT resources, and the IT requirements to improve the implementation success. Two members of the research team (PKT and RA) will conduct a 60-minute focus group in-person or virtually at each participating site with the members of the local implementation committee and IT members. We will develop the interview guide based on the existing literature on MPAI-4 implementation [23, 30, 70, 71], the team’s experience in this content area, and the questions from the CFIR interview guide tool [58, 61]. Interviews will be audio-recorded and transcribed verbatim.

**Survey:** Quantitative data will be collected from a survey sent to all clinicians. The survey will be in English or French and will include Organizational Readiness for Implementing Change (ORIC) and the implementation outcomes measures described below. Sociodemographic variables will be collected including gender, age, clinical site, profession and number years of practice. Participants will need approximately 20 minutes to complete the survey.

***Organizational Readiness for Implementing Change***

The ORIC is based on Weiner’s organizational theory [72], and aims to evaluate the organization’s readiness to implement change, including its commitment to change and its change efficacy as perceived by its members. It will also be used to guide the clinicians in identifying the strategies and resources relevant to their context [73]. The survey will contain 10 items scored on a 5-point Likert scale, ranging

from “Disagree” to “Agree” [74]. It was translated into many languages including Canadian French where the validation process has been conducted in a rehabilitation setting [74]. French and English versions of this tool have good psychometric properties including content validity, construct validity, and reliability [74]. We will use both English and French versions in the study to offer clinicians the opportunity to use the tool in their preferred language.

***Implementation outcomes: Acceptability of Intervention Measure (AIM), Intervention Appropriateness Measure (IAM), and Feasibility of Intervention Measure (FIM)***

These are three brief, validated, pragmatic and reliable measures developed by Weiner and collaborators and related to Proctor et al.’s IOF [62, 75]. Each measure has four items per construct with ordinal five response options (from “completely disagree” to “completely agree”), for a total of 12 questions [75]. Cut-off scores for interpretation are not yet available; however, we will consider the mean value as in many other studies using these measures [76, 77]. As a result, the higher scores will indicate greater acceptability, appropriateness, or feasibility [75]. AIM, IAM and FIM have satisfactory psychometric properties including good inter-item reliability and test-retest reliability [75]. We will use the original English version and a non-validated French translation of these measures because there is no French version of this tool.

**Data Analysis:** Two members of the research team (PKT and RA) will conduct a content analysis of the qualitative data [78] using N\*Vivo software (version 12) [79] and a quantitative data analysis using SAS version 12.1 [80]. They will anonymize the transcripts and review them for accuracy. Analysis will involve three phases [78]: 1) familiarizing with the data; 2) organizing the data with a categorization matrix; and, 3) reporting the data with the presentation of the described contents (meanings) of the categories and themes. Throughout this process, PKT and RA will meet regularly with one another and

1  
2 with the larger research team to discuss coding and potential recurring ideas and generate categories and  
3  
4 themes as a part of a reflexive cycle. Any discrepancies will be resolved by discussion. Trustworthiness  
5  
6 [81-83] will be supported through triangulation of data sources and collection methods (focus groups,  
7  
8 meetings minutes, facilitators' notes, and surveys), using of multiple coders and several sites. Descriptive  
9  
10 analyses of the quantitative data (e.g., mean, standard deviation, percentage) will be conducted to address  
11  
12 variation in clinician readiness. We will explore the differences between sociodemographic variables of  
13  
14 clinician readiness within and across sites. Qualitative and quantitative data will be integrated to further  
15  
16 inform the analysis as well as during the interpretation phase. We will use quotes to illustrate the most  
17  
18 relevant findings from the qualitative data. We will generate tables with information on socio-  
19  
20 demographic characteristics and from measurement tools.  
21  
22  
23  
24  
25  
26

27 **Anticipated Results:** Results from this mixed-methods design within the multiple case study will help  
28  
29 to: 1) shed light on clinicians' perceptions regarding the MPAI-4 and its compatibility with the clinical  
30  
31 practice; and 2) inform and build a multicomponent implementation blueprint, or plan of necessary  
32  
33 resources tailored to the local contexts and to improve the implementation success. The identification of  
34  
35 the barriers and enablers informed by CFIR will guide the tailoring of the implementation strategies [60].  
36  
37 We will consider implementation strategies from clinical teams as well as from ERIC after matching  
38  
39 with the identified barriers and enablers. These strategies will be ranked and prioritize by the research  
40  
41 team and the local implementation committees, to identify the strategies most likely to increase the  
42  
43 clinicians' adoption of the MPAI-4 and to inform the next steps. The variation across the sites is expected  
44  
45 to result in different adaptations to the MPAI-4 and/or implementation strategies to improve  
46  
47 implementation success. Building on existing MPAI-4 materials used during the previous  
48  
49 implementation of the MPAI-4 in traumatic brain injury rehabilitation settings [30, 51], we will develop  
50  
51 the administration protocol of the MPAI-4 in each participating site (who, what, when, where and how).  
52  
53  
54  
55  
56  
57  
58  
59  
60

### **Implementation phase**

This phase will include two components: 1) a pilot period when each clinician administers the MPAI-4 to four/six patients per site over a time selected by each site, followed by adaptations based on clinicians' and managers' feedback; and 2) a main study implementation of the MPAI-4 across all sites (objectives 3, 4 & 5). We will identify the barriers and enablers to the use of the MPAI-4 by clinical teams and evaluate the implementation outcomes. A summary of outcomes is available in Table 2. This phase is informed by the three theoretical frameworks as CFIR, IOF, and NPT.

**Table 2: Process measures of implementation evaluation**

Process results and measures	Method of data collection	Data analysis
Description of barriers and enablers of MPAI-4' implementation	Meetings minutes from local needs assessment External and internal support field notes and site journal,  Focus groups with clinicians at 6 months (CFIR)	Content analysis
Description of the experiences with MPAI-4' implementation	Focus groups with clinicians and managers at 15 months	Content analysis
Description of the degree of normalization	Surveys (NoMAD) at 6, 12 and 15 months	Descriptive statistics
Feasibility:  1. Number of clinicians who used MPAI-4 and its subscales 2. Number of clinicians who interpret MPAI-4 and its subscales 3. Time taken to complete MPAI-4 and all its subscales 4. Time required to ACCESS/Redcap platform to a report 5. Numbers of reports downloaded and put in the patient charts	Access data/Patient charts Self-report by clinicians Field notes from external and internal support	Quantitative: Descriptive statistics  Qualitative: content analysis
Fidelity:  1. Number of missing subscales by clinicians 2. Number of missing case report forms 3. Reasons for missing data	Access data/Patient charts Field notes from internal and external support	Quantitative: Descriptive statistics  Qualitative: content analysis
Acceptability:  1. % clinicians reporting MPAI-4 helpful in discussing symptoms/symptom management 2. % clinicians reporting ease of use and comprehensibility for MPAI-4 and technology systems	Surveys and self-report by clinicians Field notes form internal and external support	Quantitative: Descriptive statistics  Qualitative: content analysis



**Table 2: Process measures of implementation evaluation - Continued**

Outcomes measures	Method of data collection	Data analysis
<p>Appropriateness:</p> <ol style="list-style-type: none"> <li>1. % clinicians who find that MPAI-4 fit with workflow and clinical processes (e.g., MPAI-4-meaningful for clinical care, integrated in electronic health record system, linked clinical decision support)</li> <li>2. MPAI-4 fit with clinic culture and values</li> <li>3. The perceived relative advantage to use MPAI-4 (technology comfort, literacy level, meaningful for clinical condition)</li> </ol>	<p>Survey and self-reported by clinicians</p> <p>Field notes from external and internal support</p>	<p>Quantitative: descriptive statistics</p> <p>Qualitative: content analysis</p>
<p>Fidelity:</p> <ol style="list-style-type: none"> <li>1. Proportion of patients whose MPAI-4 was completed on expected times (admission, discharge, both)</li> <li>2. % of targeted patients whose MPAI-4 was completed at the first individual intervention plan % of targeted patients whose MPAI-4 was completed at discharge</li> <li>3. Proportion of patients whose MPAI-4 scores were sent to the follow-up team, e.g., return-to-work program or CLSC</li> <li>4. Proportion of targeted patients for whom the MPAI-4 was completed at discharge, so the total score decreased (improvement, minimal important difference and robust important difference).</li> </ol>	<p>Access data/Patient charts</p>	<p>Quantitative: descriptive analysis</p>
<p>Adoption:</p> <p>Number of clinicians who used MPAI-4 over time (6, 12 and 15 months post training)</p>	<p>Access data/Patient charts</p>	<p>Quantitative: descriptive analytics</p>
<p>Experiences/satisfaction/success</p> <p>MPAI-4 informs the individual intervention plan even though it is not discussed with the patients</p>	<p>Focus groups</p> <p>Field notes</p>	<p>Qualitative: content analysis</p>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Design:** A qualitative descriptive approach will be used to identify the barriers and enablers, and the experiences of using MPAI-4. The level of integration of the MPAI-4 in the practices and the implementation outcomes will be determined by using a quantitative component.

**Participants:** All managers and clinicians from each site will be invited to participate via email. We aim to recruit at least eight to ten clinicians, and one and/or two managers from each site to participate in focus groups. We will invite all the clinicians to fill out the survey.

**Data collection:** We will use three data collection methods: i) focus groups; ii) patients' charts; and iii) surveys. Data collection will occur at three time points: at 6-, 12-, and 15- months following the pilot period defined in each site, during the main study implementation phase. The collected data will include information about: i) barriers and enablers to the use of MPAI-4; ii) integration of the MPAI-4 into clinical practice; ii) implementation outcomes (acceptability, appropriateness, feasibility, fidelity and adoption); and iii) experiences of the MPAI-4 use (for instance: communication within and across sites, predefined goal as organizational, reflexivity, patient-centered care, patient-outcomes, technical and technological issues).

**Focus group** questions will be used to identify the factors that influence the use of the MPAI-4. We will conduct two focus groups of 90-minutes each [69]: the first, to identify the barriers and enablers with the clinicians, and the second, to explore managers' and clinicians' experiences of using MPAI-4. We will pilot-test the interview guide with one clinician and/or one manager to ensure the clarity of questions and their comprehensiveness. The first interview guide will be composed of open-ended questions addressing

different constructs of the CFIR framework such as inner and outer settings, planning strategies, individual characteristics, and intervention characteristics.

The second interview guide will consist of open-ended questions addressing clinicians and managers' viewpoints of their experiences with the use of MPAI-4 during this period, the strategies deployed for its implementation, the level of integration of the MPAI into the patient's treatment plan, their level of satisfaction, and the lessons learned. Managers and clinicians will be recruited by email by a member of the research team. Before the focus group, participants will be asked to complete a brief sociodemographic questionnaire. We will use a similar approach for the focus group in the qualitative arm for the pre-implementation phase.

**Patients' charts:** We will report MPAI-4 data from patient charts at two periods of administration (admission and discharge). Several implementation process outcomes such as fidelity, feasibility and adoption will be collected from patients' charts. For instance, we will collect data on the number/percentage of clinicians who have used the MPAI-4 at the admission, the discharge or both; all its subscales; the missing subscales; the number of clients with whom the clinicians used the MPAI-4 (See Table 2).

**Survey:** We will use a 15-minute online, auto-administrated, and anonymous survey administered to all clinicians at each of the three time points to collect information on the MPAI-4 integration over time [63, 84]. The survey includes three parts. Part A: sociodemographic information; Part B: Information about the current use and the outcomes of implementation efforts. Part C is comprised of 20 items on a 5-point Likert scale (1=completely agree to 5=completely disagree) from Normalisation Measure Development questionnaire (*NoMAD*) [85, 86], and assesses staff perceptions of the factors relevant to embedding the

1  
2 MPAI-4 in their clinical practice. This instrument has satisfactory psychometrics properties in French  
3  
4 and English including the construct validity and the reliability [63].  
5  
6  
7  
8  
9

10 **Data analysis:** We will analyze the quantitative and qualitative data separately. The qualitative data will  
11  
12 be used to inform the next steps in the implementation, mainly for choosing and tailoring implementation  
13  
14 strategies. Both the quantitative and qualitative data will inform the last qualitative data collection on  
15  
16 participants' experiences of the MPAI-4 use.  
17  
18  
19

20 The qualitative data will be analyzed using a hybrid deductive-inductive approach [87]. Themes will be  
21  
22 derived from the CFIR [58] while still allowing for emerging categories. Quantitative data from the  
23  
24 NoMAD will be analyzed using descriptive analyses (e.g., mean, standard deviation, percentage).  
25  
26 Internal consistency of all theoretical measures will be evaluated using Cronbach's alpha with an  
27  
28 acceptable threshold defined as 0.70 [88]. We will estimate the MPAI-4 subscale scores, as well as the  
29  
30 overall score by data from patients' charts. We will generate the user progress reports for each site, and  
31  
32 across sites. We will generate tables or figures with information on clinicians' socio-demographic  
33  
34 characteristics. We will use quotes to illustrate the most relevant findings.  
35  
36  
37

38 We will subsequently aggregate and integrate the data for a more comprehensive interpretation of the  
39  
40 results. Triangulation across various data sources, theoretical applications, data collection and data  
41  
42 analysis techniques will be used to increase the depth of understanding of the barriers and enablers to the  
43  
44 use of the MPAI-4 in stroke rehabilitation settings. We will highlight all areas of convergence and  
45  
46 divergence across the participating rehabilitation sites.  
47  
48  
49  
50  
51  
52  
53

54 **Anticipated Results:** This phase will generate a list of identified barriers and enablers, and potential  
55  
56 tailored implementation strategies matched to reduce the barriers, and enhance and maintain the enablers  
57  
58  
59  
60

[60]. The data will help to understand: current MPAI-4 use and the likelihood of using the MPAI-4 in the future, clinicians' perceptions of the impact of MPAI-4 on their clinical practice, the perceived changes over the time, and whether it could become a routine part of practice.

Timeline: This study was started in June 2020 and is estimated to end in December 2023. A detailed timeline is provided in Figure 1.

## DISCUSSION

We propose that successful implementation of the MPAI-4 requires an iKT approach with active engagement from all stakeholders, including managers and clinicians in each of the participating clinical rehabilitation programs. This study has the potential to contribute to the advancement of the knowledge regarding the implementation of the MPAI-4 by examining similarities and differences between sites, highlighting the influence of context in the success of implementation (or not) that may influence the providers' experiences. The combination of multiple theoretical frameworks offers the opportunity to map the broad barriers and enablers influencing clinicians' adoption of MPAI-4, and to select tailored strategies and to comprehensively measure the implementation processes and outcomes, including clinician's perceived clinical utility of MPAI-4 [70], as well as, strategies that will increase the success of the MPAI-4's implementation in the stroke rehabilitation programs.

The implementation process of the MPAI-4 will provide us with data to facilitate communication between clinical teams across the care continuum. It will strengthen patient-centered practice and improve patient outcomes with the goal of optimizing social participation. This study will enhance the clinical utility of the MPAI-4 in stroke. Ultimately, we will conduct a comparison of our results on determinants with those of previous studies conducted with traumatic brain injury patients [22, 30, 89].

1  
2 Results of this study will provide insights about iKT approach within stroke rehabilitation settings that  
3  
4 could be applied to future research projects.  
5

6  
7 Despite the valuable insights that may be gained from this study, there are some limitations. Mixed  
8  
9 methods are useful to address complex research problems that require several data sources such as self-  
10  
11 reported questionnaires, patient charts, interviews and focus groups. However, self-reported data  
12  
13 collection methods and focus groups may introduce response bias, either as an under or an overestimation  
14  
15 of the expected behavior and social desirability. To overcome these challenges and potential biases, we  
16  
17 will use triangulation across data sources, various theoretical applications, different methods of analysis  
18  
19 and clinical teams' involvement to increase the in-depth understanding of our data and mitigate their  
20  
21 potential impacts [90]. To overcome these challenges/bias, we will use triangulation across data sources,  
22  
23 various theoretical applications, different methods of analysis and clinical teams' involvement to increase  
24  
25 the in-depth understanding of our data and mitigate their potential impacts.  
26  
27  
28  
29  
30  
31  
32  
33

34 **Ethics and dissemination**

35  
36  
37 The project has received the Institutional Review Board (IRB) approval of CRIR (CRIR 1523-0221/MP-  
38  
39 50-2022-968). Any protocol modifications will be submitted for approval to the IRB, prior to the  
40  
41 implementation. The project will be conducted from 2021 to 2024.  
42  
43

44  
45 The dissemination plan will be developed with the implementation team in each site. The iKT approach  
46  
47 (participatory approach, reflective approach, local team) will allow for ongoing knowledge exchange  
48  
49 within and between participating sites. In addition, we will disseminate results in peer-reviewed  
50  
51 publications and at local, national, and international scientific conferences/workshops. Reporting of this  
52  
53 study will seek to satisfy the standards of Good Reporting of a Mixed Methods Study [91] and the  
54  
55 Standards for Reporting Implementation Studies [92, 93].  
56  
57  
58  
59  
60

**Acknowledgements:** We would like to acknowledge the leadership of managers in the participating rehabilitation centers, and we thank the members of the various local implementation committees and all the clinicians in these sites.

**Authors' contributions:** PKT led the writing of the manuscript. PKT, SA and AT led the study design and produced the first draft of the article including all sections. CA, MM, WW, FP and RA contributed substantially to the study design and provided critical feedback on the manuscript. All authors read and approved the final version of the manuscript.

**Funding statement:** This study has received a grant from the *Pôle Universitaire en Réadaptation* (PUR) of the CRIR grant number PUR 2018-2019, and salary awards from the Fonds de recherche du Québec – Santé (FRQS) for SA, AT, CA, WW. Funding was obtained by SA and AT to support the proposed research. This funding has been used to give students awards (PKT and RA). The implementation of the MPAI-4 is supported by the Biomedical Research and Informatics Living Laboratory for Innovative Advances of New Technologies (BRILLIANT) in Community Mobility Rehabilitation program funded by the Canadian Foundation of Innovation and the Ministry of Health of Quebec (#36053). CRIR is funded by the Fonds de recherche du Québec – Santé, et Société et culture.

**Competing interests' statement:** The authors declare no competing interests.

**Authors' information:**

1. School of Physical and Occupational Therapy, McGill University, Montréal, Canada
2. Integrated University Health and Social Services Centre for West-Central Montreal (CIUSSS du Centre-Ouest-de-l'Île-de-Montréal)
3. Integrated University Health and Social Services Centre for South-Central Montreal (Institut universitaire sur la réadaptation en déficience physique de Montréal, CIUSSS du Centre-Sud-de-l'Île-de-Montréal)

4. Centre for Interdisciplinary Research in Rehabilitation of Greater Montreal

5. Integrated Health and Social Services Centre of Laval (CISSS de Laval)

6. Institute of Health Sciences Education, McGill University, Montréal, Canada

7. School of Rehabilitation, Université de Montréal, Montréal, Canada

8. Department of Psychology, Université de Montréal, Montréal, Canada

9. School of Optometry, Université de Montréal, Montréal, Canada

10. Institut Nazareth et Louis-Braille, Integrated Health and Social Services Centre of  
Montérégie-Centre (CISSS de la Montérégie-Centre)



## REFERENCES

1. Lindsay MP, Norrving B, Sacco RL, Brainin M, Hacke W, Martins S, et al. World Stroke Organization (WSO): Global Stroke Fact Sheet 2019. *International Journal of Stroke*. 2019;14(8):806-17.
2. World stroke organization (WSO). Global Stroke Fact Sheet 2019 [Available from: [https://www.world-stroke.org/assets/downloads/WSO\\_Global\\_Stroke\\_Fact\\_Sheet.pdf](https://www.world-stroke.org/assets/downloads/WSO_Global_Stroke_Fact_Sheet.pdf)].
3. Jolliffe L, Lannin NA, Cadilhac DA, Hoffmann T. Systematic review of clinical practice guidelines to identify recommendations for rehabilitation after stroke and other acquired brain injuries. *BMJ open*. 2018;8(2):e018791.
4. Johnson CO, Nguyen M, Roth GA, Nichols E, Alam T, Abate D, et al. Global, regional, and national burden of stroke, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet Neurology*. 2019;18(5):439-58.
5. Avan A, Digaleh H, Di Napoli M, Stranges S, Behrouz R, Shojaeianbabaei G, et al. Socioeconomic status and stroke incidence, prevalence, mortality, and worldwide burden: an ecological analysis from the Global Burden of Disease Study 2017. *BMC Medicine*. 2019;17(1):191.
6. Global burden diseases Risk Factor Collaborators. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392(10159):1923-94.
7. Institute for Health Metrics and Evaluation (IHME). Findings from the Global Burden of Disease Study 2017. Seattle, WA: IHME. 2018 Accessed June 4, 2021.
8. Health and Stroke Foundation Canadian partnerships for stroke recovery. Background concepts in stroke rehabilitation. Canada; 2018. Accessed June 3, 2021.

1  
2 9. Ferro JM, Caeiro L, Figueira ML. Neuropsychiatric sequelae of stroke. Nature reviews  
3  
4 Neurology. 2016;12(5):269-80.  
5  
6 10. Zorowitz RD, Gillard PJ, Brainin M. Poststroke spasticity: sequelae and burden on stroke  
7  
8 survivors and caregivers. Neurology. 2013;80(3 Suppl 2):S45-52.  
9  
10 11. Rowe FJ, Hepworth LR, Howard C, Hanna KL, Cheyne CP, Currie J. High incidence and  
11  
12 prevalence of visual problems after acute stroke: An epidemiology study with implications for service  
13  
14 delivery. PloS one. 2019;14(3):e0213035.  
15  
16 12. Fondation des maladies du coeur et de l'AVC. Recommandations canadiennes pour les pratiques  
17  
18 optimales de soins de l'AVC. Réadaptation, rétablissement et participation communautaire  
19  
20 après un AVC: Deuxième partie : Les transitions et la participation communautaire après un AVC.  
21  
22 Canada,; 2019.  
23  
24 13. Ministère de la Santé et des Services sociaux (MSSS). Continuum de services pour les personnes  
25  
26 à risque de subir ou ayant subi un accident vasculaire cérébral: Plan de mise en œuvre Phase 2016-2018.  
27  
28 Québec; 2016.  
29  
30 14. Cieza A. Rehabilitation the Health Strategy of the 21st Century, Really? Archives of physical  
31  
32 medicine and rehabilitation. 2019;100(11):2212-4.  
33  
34 15. Negrini S, Mills J-A, Arienti C, Kiekens C, Cieza A. "Rehabilitation Research Framework for  
35  
36 Patients With COVID-19" Defined by Cochrane Rehabilitation and the World Health Organization  
37  
38 Rehabilitation Programme. Archives of physical medicine and rehabilitation. 2021;102(7):1424-30.  
39  
40 16. Krug E, Cieza A. Strengthening health systems to provide rehabilitation services. Bull World  
41  
42 Health Organ. 2017;95(3):167.  
43  
44 17. Bernhardt J, Hayward KS, Kwakkel G, Ward NS, Wolf SL, Borschmann K, et al. Agreed  
45  
46 definitions and a shared vision for new standards in stroke recovery research: The Stroke Recovery and  
47  
48 Rehabilitation Roundtable taskforce. International Journal of Stroke. 2017;12(5):444-50.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

18. World Health Organization (WHO). Rehabilitation 2030 Initiative <https://www.who.int/initiatives/rehabilitation-2030> Geneva 2017.
19. Pollock A, Baer G, Campbell P, Choo PL, Forster A, Morris J, et al. Physical rehabilitation approaches for the recovery of function and mobility following stroke. The Cochrane database of systematic reviews. 2014;2014(4):Cd001920.
20. Fricke J. Measuring Outcomes in Rehabilitation: A Review. The British Journal of Occupational Therapy. 1993;56:217 - 21.
21. Harris K, Dawson J, Gibbons E, Lim CR, Beard DJ, Fitzpatrick R, et al. Systematic review of measurement properties of patient-reported outcome measures used in patients undergoing hip and knee arthroplasty. Patient Relat Outcome Meas. 2016;7:101-8.
22. Institut national d'excellence en santé et en services sociaux (INESSS). Trousse de soutien à l'implantation clinique du MPAI-4. Québec, Québec; 2019.
23. Malec JF. The Mayo-Portland Adaptability Inventory. The Center for Outcome Measurement in Brain Injury. <http://www.tbims.org/combi/mpai> 2005.
24. Colquhoun HL, Lamontagne M-E, Duncan EA, Fiander M, Champagne C, Grimshaw JM. A systematic review of interventions to increase the use of standardized outcome measures by rehabilitation professionals. Clinical Rehabilitation. 2017;31(3):299-309.
25. Sullivan JE, Crowner BE, Kluding PM, Nichols D, Rose DK, Yoshida R, et al. Outcome Measures for Individuals With Stroke: Process and Recommendations From the American Physical Therapy Association Neurology Section Task Force. Physical Therapy. 2013;93(10):1383-96.
26. Winstein CJ, Stein J, Arena R, Bates B, Cherney LR, Cramer SC, et al. Guidelines for Adult Stroke Rehabilitation and Recovery. Stroke. 2016;47(6):e98-e169.
27. Bellon K, Malec JF, Kolakowsky-Hayner SA. Mayo-Portland Adaptability Inventory-4. J Head Trauma Rehabil. 2012;27(4):314-6.

1  
2 28. Accreditation Canada. Achieving Distinction in Stroke Care 2019 [Available from:  
3  
4 <https://accreditation.ca/news/distinction-in-stroke-care/>.  
5  
6  
7 29. Lageman SK. Commission on Accreditation of Rehabilitation Facilities (CARF). In: Kreutzer JS,  
8 DeLuca J, Caplan B, editors. Encyclopedia of Clinical Neuropsychology. New York, NY: Springer New  
9 York; 2011. p. 647-50.  
10  
11  
12  
13 30. Guerrette MC, McKerral M. Validation of the Mayo-Portland Adaptability Inventory-4 (MPAI-  
14 4) and reference norms in a French-Canadian population with traumatic brain injury receiving  
15 rehabilitation. Disability and rehabilitation. 2021;1-7 DOI:.1080/09638288.2021.1924882.  
16  
17  
18  
19  
20 31. Malec JF, Kragness M, Evans RW, Finlay KL, Kent A, Lezak MD. Further psychometric  
21 evaluation and revision of the Mayo-Portland Adaptability Inventory in a national sample. The Journal  
22 of head trauma rehabilitation. 2003;18(6):479-92.  
23  
24  
25  
26  
27 32. Malec JF. Comparability of Mayo-Portland Adaptability Inventory ratings by staff, significant  
28 others and people with acquired brain injury. Brain injury. 2004;18(6):563-75.  
29  
30  
31  
32 33. Williams E, Jackson H, Wagland J, Martini A. Community Rehabilitation Outcomes for Different  
33 Stroke Diagnoses: An Observational Cohort Study. Archives of rehabilitation research and clinical  
34 translation. 2020;2(2):100047.  
35  
36  
37  
38  
39 34. (MSSS) MdI SedSs. Continuum de services pour les personnes à risque de subir ou ayant subi un  
40 accident vasculaire cérébral Quebec; 2018.  
41  
42  
43  
44 35. Harvey G, Loftus-Hills A, Rycroft-Malone J, Titchen A, Kitson A, McCormack B, et al. Getting  
45 evidence into practice: the role and function of facilitation. Journal of Advanced Nursing.  
46 2002;37(6):577-88.  
47  
48  
49  
50 36. Kitson A, Harvey G, McCormack B. Enabling the implementation of evidence based practice: a  
51 conceptual framework. Quality in Health Care. 1998;7(3):149-58.  
52  
53  
54  
55  
56  
57  
58  
59  
60

37. Damschroder LJ. Clarity out of chaos: Use of theory in implementation research. *Psychiatry Research*. 2020;283:112461.
38. Powell BJ, Waltz TJ, Chinman MJ, Damschroder LJ, Smith JL, Matthieu MM, et al. A refined compilation of implementation strategies: results from the Expert Recommendations for Implementing Change (ERIC) project. *Implementation Science*. 2015;10(1):21.
39. Bodenheimer T, Sinsky C. From triple to quadruple aim: care of the patient requires care of the provider. *Ann Fam Med*. 2014;12(6):573-6.
40. Sikka R, Morath JM, Leape L. The Quadruple Aim: care, health, cost and meaning in work. *BMJ Quality & Safety*. 2015;24(10):608.
41. Kothari A, McCutcheon C, Graham ID. Defining Integrated Knowledge Translation and Moving Forward: A Response to Recent Commentaries. *Int J Health Policy Manag*. 2017;6(5):299-300.
42. Graham ID, Kothari A, McCutcheon C, Angus D, Banner D, Bucknall T, et al. Moving knowledge into action for more effective practice, programmes and policy: protocol for a research programme on integrated knowledge translation. *Implementation Science*. 2018;13(1):22.
43. Thomas G. A Typology for the Case Study in Social Science Following a Review of Definition, Discourse, and Structure. *Qualitative Inquiry*. 2011;17(6):511-21.
44. Eisenhardt KM, Graebner ME. Theory Building From Cases: Opportunities And Challenges. *Academy of Management Journal*. 2007;50(1):25-32.
45. Yin RK. *Case study research: Design and methods* (4th Ed.). Thousand Oaks, CA: Sage. 2009.
46. Yin RK. *Case Study Research and Applications: Design and Methods* (6th ed.). Thousand Oaks, CA: Sage 2018.
47. Creswell JW. *Research Design: Qualitative, Quantitative, and mixed methods approaches*. California: Sage Publication; 2009.

1  
2 48. Palinkas LA, Aarons GA, Horwitz S, Chamberlain P, Hurlburt M, Landsverk J. Mixed method  
3  
4 designs in implementation research. *Administration and policy in mental health*. 2011;38(1):44-53.  
5  
6 49. Hamed R, Tariah HA, Malkawi S, Holm MB. The Arabic version of the Mayo-Portland  
7  
8 Adaptability Inventory 4: a validation study. *International journal of rehabilitation research*  
9  
10  
11 *Internationale Zeitschrift fur Rehabilitationsforschung Revue internationale de recherches de*  
12  
13 *readaptation*. 2012;35(3):243-7.  
14  
15 50. Cattelani R, Corsini D, Posteraro L, Agosti M, Saccavini M. The Italian version of the Mayo-  
16  
17 Portland Adaptability Inventory-4. A new measure of brain injury outcome. *European journal of physical*  
18  
19 *and rehabilitation medicine*. 2009;45(4):513-9.  
20  
21 51. McKerral M, Léveillé G, Goulet P, Vincent P. Manuel MPAI-4 Version canadienne-française.  
22  
23 The Center for Outcome Measurement in Brain Injury. Available from:  
24  
25 <http://www.tbims.org/combi/mpai/>. 2014 [Accessed June 6, 2022].  
26  
27 52. Ataman R TA, Roberge-Dao J, et al. . Measurement Properties of the Mayo-Portland Adaptability  
28  
29 Inventory (MPAI-4) and related measures: A Systematic Review [published online ahead of print, 2023  
30  
31 Jan 25]. . *Arch Phys Med Rehabil*. 2023:S0003-9993(23)00046-1.  
32  
33 53. Jackson D, Seaman K, Sharp K, Singer R, Wagland J, Turner-Stokes L. Staged residential post-  
34  
35 acute rehabilitation for adults following acquired brain injury: A comparison of functional gains rated on  
36  
37 the UK Functional Assessment Measure (UK FIM+FAM) and the Mayo-Portland Adaptability Inventory  
38  
39 (MPAI-4). *Brain Inj*. 2017;31(11):1405-13.  
40  
41 54. Nayar M, Vanderstay R, Siegert RJ, Turner-Stokes L. The UK Functional Assessment Measure  
42  
43 (UK FIM+FAM): Psychometric Evaluation in Patients Undergoing Specialist Rehabilitation following a  
44  
45 Stroke from the National UK Clinical Dataset. *PloS one*. 2016;11(1):e0147288.  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

55. Altman IM, Swick S, Malec JF. Effectiveness of Home- and Community-Based Rehabilitation in a Large Cohort of Patients Disabled by Cerebrovascular Accident: Evidence of a Dose-Response Relationship. *Archives of Physical Medicine and Rehabilitation*. 2013;94(9):1837-41.
56. Malec JF. Malec, J.F. The Mayo-Portland Adaptability Inventory. The Center for Outcome Measurement in Brain Injury. <http://www.tbims.org/combi/mpai>. 2005 [Accessed January 3, 2021].
57. Kilbourne AM, Goodrich DE, Miake-Lye I, Braganza MZ, Bowersox NW. Quality Enhancement Research Initiative Implementation Roadmap: Toward Sustainability of Evidence-based Practices in a Learning Health System. *Medical care*. 2019;57.
58. Damschroder LJ, Aron DC, Keith RE, Kirsh SR, Alexander JA, Lowery JC. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implementation Science*. 2009;4(1):50.
59. Esmail R, Hanson HM, Holroyd-Leduc J, Brown S, Strifler L, Straus SE, et al. A scoping review of full-spectrum knowledge translation theories, models, and frameworks. *Implementation Science*. 2020;15(1):11.
60. Waltz TJ, Powell BJ, Fernández ME, Abadie B, Damschroder LJ. Choosing implementation strategies to address contextual barriers: diversity in recommendations and future directions. *Implementation science : IS*. 2019;14(1):42.
61. Consolidated Framework for Implementation research (CFIR). Welcome to the Interview Guide Tool: CFIR Domains [https://cfirguide.org/guide/app/-/guide\\_select](https://cfirguide.org/guide/app/-/guide_select). [Accessed January 12, 2021].
62. Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A, et al. Outcomes for implementation research: conceptual distinctions, measurement challenges, and research agenda. *Administration and policy in mental health*. 2011;38(2):65-76.

1  
2 63. Finch TL, Girling M, May CR, Mair FS, Murray E, Treweek S, et al. Improving the normalization  
3  
4 of complex interventions: part 2 - validation of the NoMAD instrument for assessing implementation  
5  
6 work based on normalization process theory (NPT). BMC Medical Research Methodology.  
7  
8 2018;18(1):135.  
9  
10  
11 64. Finch TL, Rapley T, Girling M, Mair FS, Murray E, Treweek S, et al. Improving the  
12  
13 normalization of complex interventions: measure development based on normalization process theory  
14  
15 (NoMAD): study protocol. Implementation Science. 2013;8(1):43.  
16  
17  
18 65. Rapley T, Girling M, Mair FS, Murray E, Treweek S, McColl E, et al. Improving the  
19  
20 normalization of complex interventions: part 1 - development of the NoMAD instrument for assessing  
21  
22 implementation work based on normalization process theory (NPT). BMC Med Res Methodol.  
23  
24 2018;18(1):133.  
25  
26  
27 66. Vis C, Ruwaard J, Finch TL, Rapley T, de Beurs D, van Stel H, et al. Toward an Objective  
28  
29 Assessment of Implementation Processes for Innovations in Health Care: Psychometric Evaluation of  
30  
31 the Normalization Measure Development (NoMAD) Questionnaire Among Mental Health Care  
32  
33 Professionals. J Med Internet Res. 2019;21(2):e12376.  
34  
35  
36 67. Lewis CC, Boyd MR, Walsh-Bailey C, Lyon AR, Beidas R, Mittman B, et al. A systematic review  
37  
38 of empirical studies examining mechanisms of implementation in health. Implementation Science.  
39  
40 2020;15(1):21.  
41  
42  
43 68. Carpenter C. Using qualitative focus groups to evaluate health programmes and service delivery.  
44  
45 Qualitative research in evidence-based rehabilitation: Elsevier; 2004. p. 51-64.  
46  
47  
48 69. Krueger RA, Casey MA. Focus groups : a practical guide for applied research: Thousand Oaks,  
49  
50 California : SAGE; 2009.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2 70. Kean J, Malec JF, Cooper DB, Bowles AO. Utility of the Mayo-Portland adaptability inventory-4  
3  
4 for self-reported outcomes in a military sample with traumatic brain injury. Archives of physical  
5  
6 medicine and rehabilitation. 2013;94(12):2417-24.  
7  
8  
9 71. Wilde EA, Whiteneck GG, Bogner J, Bushnik T, Cifu DX, Dikmen S, et al. Recommendations  
10  
11 for the use of common outcome measures in traumatic brain injury research. Archives of physical  
12  
13 medicine and rehabilitation. 2010;91(11):1650-60.e17.  
14  
15  
16 72. Weiner BJ. A theory of organizational readiness for change. Implementation Science.  
17  
18 2009;4(1):67.  
19  
20 73. Shea CM, Jacobs SR, Esserman DA, Bruce K, Weiner BJ. Organizational readiness for  
21  
22 implementing change: a psychometric assessment of a new measure. Implementation Science.  
23  
24 2014;9(1):7.  
25  
26  
27 74. Ruest M, Léonard G, Thomas A, Desrosiers J, Guay M. French cross-cultural adaptation of the  
28  
29 Organizational Readiness for Implementing Change (ORIC). BMC Health Services Research.  
30  
31 2019;19(1):535.  
32  
33  
34 75. Weiner BJ, Lewis CC, Stanick C, Powell BJ, Dorsey CN, Clary AS, et al. Psychometric  
35  
36 assessment of three newly developed implementation outcome measures. Implementation Science.  
37  
38 2017;12(1):108.  
39  
40  
41 76. Pellet J, Weiss M, Zúñiga F, Mabire C. Implementation and preliminary testing of a theory-  
42  
43 guided nursing discharge teaching intervention for adult inpatients aged 50 and over with multimorbidity:  
44  
45 a pragmatic feasibility study protocol. Pilot and Feasibility Studies. 2021;7(1).  
46  
47  
48 77. Eisman AB, Palinkas, L. A., Brown, S., Lundahl, L., & Kilbourne, A. M. . A mixed methods  
49  
50 investigation of implementation determinants for a school-based universal prevention intervention. .  
51  
52 Implementation Research and Practice 2022;3.  
53  
54  
55 78. Elo S, Kyngäs H. The qualitative content analysis process. J Adv Nurs. 2008;62(1):107-15.  
56  
57  
58  
59  
60

1  
2 79. 12 N. Software in Qualitative Data Analysis: QSR International 2020 [Available from:  
3  
4 [https://www.qsrinternational.com/nvivo-qualitative-data-analysis-software/support-services/customer-](https://www.qsrinternational.com/nvivo-qualitative-data-analysis-software/support-services/customer-hub/research-ready-(1)/research-ready)  
5  
6 [hub/research-ready-\(1\)/research-ready](https://www.qsrinternational.com/nvivo-qualitative-data-analysis-software/support-services/customer-hub/research-ready-(1)/research-ready).  
7  
8  
9 80. SAS Institute Inc. SAS. Version 12.1.  
10  
11 <https://support.sas.com/documentation/onlinedoc/stat/121/intro.pdf>. 2020.  
12  
13 81. Shenton AK. Strategies for Ensuring Trustworthiness in Qualitative Research Projects. Education  
14 for Information. 2004;22:63-75.  
15  
16 82. Graneheim UH, Lundman B. Qualitative content analysis in nursing research: concepts,  
17 procedures and measures to achieve trustworthiness. Nurse education today. 2004;24(2):105-12.  
18  
19 83. Elo S, Kääriäinen M, Kanste O, Pölkki T, Utriainen K, Kyngäs H. Qualitative Content Analysis:  
20 A Focus on Trustworthiness. SAGE Open. 2014;4(1):2158244014522633.  
21  
22 84. ImpleMentAll. Normalization Measure Development Questionnaire (NoMAD), (2018, October  
23 16). Retrieved from <http://www.implementall.eu/> 2018  
24  
25 85. Finch TL, Girling M, May CR, Mair FS, Murray E, Treweek S, et al. Nomad: Implementation  
26 measure based on Normalization Process Theory. [Measurement instrument]. Retrieved from  
27 <http://www.normalizationprocess.org>. 2015.  
28  
29 86. May CR, Cummings A, Girling M, Bracher M, Mair FS, May CM, et al. Using Normalization  
30 Process Theory in feasibility studies and process evaluations of complex healthcare interventions: a  
31 systematic review. Implementation Science. 2018;13(1):80.  
32  
33 87. Hsieh H-F, Shannon S. Three Approaches to Qualitative Content Analysis. Qualitative health  
34 research. 2005;15:1277-88.  
35  
36 88. Cronbach LJ. Coefficient alpha and the internal structure of tests. Psychometrika. 1951;16:297-  
37 334.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2 89. Lamontagne M-E, Bussi res A, Kairy D, Ahmed S, Zidarov D, Messier F.  valuation des effets  
3 de l'implantation de quatre nouvelles pratiques cliniques fond es sur des donn es probantes- Phase 2  
4 2017.  
5  
6  
7  
8 90. Joseph W Sakshaug AC, Trivellore E Raghunathan, . Do Sequential Mixed-Mode Surveys  
9 Decrease Nonresponse Bias, Measurement Error Bias, and Total Bias? An Experimental Study. Journal  
10 of Survey Statistics and Methodology. 2019;7 (4):545–71.  
11  
12  
13 91. O'Cathain A, Murphy E, Nicholl J. The quality of mixed methods studies in health services  
14 research. Journal of health services research & policy. 2008;13(2):92-8.  
15  
16 92. Smith JD, Li DH, Rafferty MR. The Implementation Research Logic Model: a method for  
17 planning, executing, reporting, and synthesizing implementation projects. Implementation Science.  
18 2020;15(1):84.  
19  
20 93. Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of  
21 interventions: template for intervention description and replication (TIDieR) checklist and guide. BMJ :  
22 British Medical Journal. 2014;348:g1687.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Figure 1: Implementation design and Timeline of MPAI implementation project**

For peer review only

Figure 1: Implementation design and Timeline of MPAI-4 implementation project

